

Decay Constants of Pseudoscalar Mesons to Two Loops in Three-Flavor Partially Quenched χ PT

Johan Bijnens and Timo A. Lähde

Department of Theoretical Physics, Lund University,
Sölvegatan 14A, S 223 62 Lund, Sweden

Abstract

This paper presents a first study of the decay constants of the charged, or flavor-off-diagonal, pseudoscalar mesons to two loops for three flavors of sea quarks, in Partially Quenched Chiral Perturbation Theory (PQ χ PT). Explicit analytical expressions up to $\mathcal{O}(p^6)$ in the momentum expansion are given. The calculations have been performed within the supersymmetric formulation of PQ χ PT. We also present some numerical results to indicate the size of the corrections.

PACS: 12.38.Gc, 12.39.Fe, 11.30.Rd

Decay Constants of Pseudoscalar Mesons to Two Loops in Three-Flavor Partially Quenched χ PT

Johan Bijnens¹ and Timo A. Lähde¹

¹*Department of Theoretical Physics, Lund University,
Sölvegatan 14A, S 223 62 Lund, Sweden*

This paper presents a first study of the decay constants of the charged, or flavor-off-diagonal, pseudoscalar mesons to two loops for three flavors of sea quarks, in Partially Quenched Chiral Perturbation Theory (PQ χ PT). Explicit analytical expressions up to $\mathcal{O}(p^6)$ in the momentum expansion are given. The calculations have been performed within the supersymmetric formulation of PQ χ PT. We also present some numerical results to indicate the size of the corrections.

PACS numbers: 12.38.Gc, 12.39.Fe, 11.30.Rd

I. INTRODUCTION

The derivation of low-energy hadronic observables, e.g. meson masses and decay constants, from the theory of the strong interaction (QCD) has so far proven to be impossible by means of analytical methods. Because of this situation, numerical Lattice QCD simulations, whereby the functional integral is evaluated numerically on a discretized space-time lattice, have developed into a major field of study. Such simulations are, however, seriously hampered by difficulties in the simulation of dynamical sea quark effects. Although much progress has been made recently, it is still impractical, for computational reasons, to simulate with sea quark masses that are close to the physical u, d quark masses of a few MeV. This situation, with sea quark masses of a few tens of MeV, is referred to as partially quenched (PQ) QCD. Consequently, the physical values of the sea quark masses have to be reached by extrapolation from the partially quenched simulation results.

A practical method for this extrapolation is provided by Chiral Perturbation Theory (χ PT), which provides the correct quark mass dependences of the various physical quantities that are measured on the lattice. Standard three-flavor χ PT as introduced by Weinberg, Gasser and Leutwyler in Refs. [1], is valid in the (unquenched) QCD case of equal valence and sea quark masses. The generalization of χ PT to the quenched case (without sea quarks) or to the partially quenched case (sea quark masses different from the valence ones) has been carried out by Bernard and Golterman in Refs. [2, 3]. The quark mass dependence of partially quenched chiral perturbation theory (PQ χ PT) is explicit, and thus the limit where the sea quark masses become equal to the valence quark masses can be taken. As a consequence, χ PT is included in PQ χ PT and the free parameters, or low-energy constants (LEC:s), of χ PT can be directly obtained from those of PQ χ PT [3, 4].

The calculation of charged pseudoscalar meson masses and decay constants to one loop (NLO) in PQ χ PT has been carried out in Refs. [3, 4, 5], and first results for the mass of a charged pseudoscalar meson at two loops or next-to-next-to-leading order (NNLO) in PQ χ PT have

already appeared, for degenerate sea quark masses, in Ref. [6]. The need for such calculations is clear as NNLO effects have already been detected in Lattice QCD simulations [7, 8]. A calculation of the pseudoscalar meson masses for nondegenerate sea quarks is in progress [9].

This paper presents the first calculation of the decay constants of the charged, or flavor off-diagonal, pseudoscalar mesons in NNLO PQ χ PT, for three flavors of sea quarks ($n_{\text{sea}} = 3$). The results are characterized by the number of nondegenerate valence and sea quarks, denoted d_{val} and d_{sea} , respectively. For the decay constants of the charged pseudoscalar mesons, the maximum number of nondegenerate valence quark masses is $d_{\text{val}} = 2$. The degree of quark mass degeneracy in each result is sometimes also referred to with the notation $d_{\text{val}} + d_{\text{sea}}$. The decay constant of the charged pion in the $SU(2)$ symmetric limit thus corresponds to the $1 + 2$ case. Likewise, the decay constants of the charged and neutral kaons may be obtained from the $d_{\text{val}} = 2$ results with $d_{\text{sea}} = 2$. Results are also presented for the case of $d_{\text{sea}} = 1$ (all sea quark masses equal), and $d_{\text{sea}} = 3$ (all sea quark masses different). An extension of the present work to the neutral pseudoscalar mesons is also planned.

The analytical expressions for the NNLO shift of the decay constants are in general very long, but the expressions simplify considerably when pairs of sea or valence quark masses become degenerate. In view of this, the NNLO loop results are given separately for each case of $d_{\text{val}} + d_{\text{sea}}$ considered. In the next sections, the technical background for the NNLO calculations, the full results for the decay constants of the charged pseudoscalar mesons and numerical results as a function of the input quark masses are given, along with a concluding discussion.

II. TECHNICAL OVERVIEW

Most of the technical aspects that concern the calculation of the pseudoscalar meson decay constants to two loops, or NNLO, are identical to those of the calculation of the pseudoscalar meson mass, and have already been justified in Ref. [6]. Most significantly, the Lagrangians of PQ χ PT at $\mathcal{O}(p^4)$ and $\mathcal{O}(p^6)$ may be directly

obtained from the corresponding Lagrangians of normal (unquenched) n_F flavor χ PT, provided that the traces and meson matrices are replaced with the supertraces and meson matrices relevant to the partially quenched theory [2, 3, 4]. This can be argued from the Replica method as in Ref. [10], or by the fact that all the relations used to constrain the number of terms in Ref. [11] remain valid when traces are changed to supertraces. We work here in the version of PQ χ PT without the Φ_0 as discussed in Ref. [5].

A. Masses and Low-Energy Constants

All calculations in this paper have been performed with three flavors of valence quarks, three flavors of sea quarks and three flavors of bosonic 'ghost' quarks. These may be viewed as the u, d and s quarks in the valence, sea and ghost sectors, respectively. The purpose of the ghost quarks is to remove the valence quark loops which are disconnected from the external legs.

The input quark masses m_i enter into the calculation in terms of the lowest order squared meson masses χ_i , which are defined as usual in χ PT, by $\chi_i = 2B_0 m_i$. In the present calculations, we thus have three valence inputs χ_1, χ_2, χ_3 , three sea inputs χ_4, χ_5, χ_6 , and three ghost inputs χ_7, χ_8, χ_9 . In order for the disconnected valence quark loops to be canceled, the masses of the ghost quarks are always equal to those of the corresponding valence quarks, such that $\chi_7 = \chi_1$, $\chi_8 = \chi_2$ and $\chi_9 = \chi_3$. Explicitly, for $d_{\text{val}} = 1$, we have $\chi_1 = \chi_2 = \chi_3$, for $d_{\text{val}} = 2$, we have $\chi_1 = \chi_2 \neq \chi_3$ and for $d_{\text{val}} = 3$, we have $\chi_1 \neq \chi_2 \neq \chi_3$. Similarly, for the sea quarks $d_{\text{sea}} = 1$ implies $\chi_4 = \chi_5 = \chi_6$, while $d_{\text{sea}} = 2$ implies $\chi_4 = \chi_5 \neq \chi_6$ and finally $d_{\text{sea}} = 3$ that $\chi_4 \neq \chi_5 \neq \chi_6$.

The number of independent low-energy constants in unquenched and partially quenched χ PT is slightly different, but the former are always linear combinations of the latter. For PQ χ PT, they are F_0 and B_0 at leading order, L_0^r through L_{12}^r at NLO and K_1^r through K_{115}^r at NNLO. In contrast, for three flavor unquenched χ PT they are F_0 and B_0 at leading order, $L_0^{r(3)}$ through $L_{12}^{r(3)}$ at NLO and C_1^r through C_{94}^r at NNLO. Note that the parameters $L_{11,12}^r$ and $L_{11,12}^{r(3)}$ correspond to the usual $H_{1,2}^r$. Also, the parameters F_0 and B_0 are identical for unquenched and partially quenched χ PT. At NLO, the relations between the low-energy constants are [1, 11, 12]

$$\begin{aligned} L_1^{r(3)} &= L_0^r/2 + L_1^r, \\ L_2^{r(3)} &= L_0^r + L_2^r, \\ L_3^{r(3)} &= -2L_0^r + L_3^r, \\ L_{4\dots 12}^{r(3)} &= L_{4\dots 12}^r, \end{aligned} \quad (1)$$

and the corresponding linear relations relevant for the NNLO parameters can be found in Ref. [11].

B. Meson Propagators in PQ χ PT

The calculation of the pseudoscalar meson mass in Ref. [6] was only performed for $d_{\text{val}} = d_{\text{sea}} = 1$. However, the present calculation of the decay constants to NNLO is also concerned with the more general cases of d_{val} and $d_{\text{sea}} = 2, 3$. This leads to much more involved expressions because of the appearance of the residues of the neutral meson propagators. We recall here the results of Ref. [5] which we have translated to Minkowski space from the Euclidean formalism used there.

In Minkowski space, the propagator G_{ij}^c of a charged, or flavor-off-diagonal meson with flavor structure $q_i \bar{q}_j$ in supersymmetric PQ χ PT, is given by [5]

$$-i G_{ij}^c(k) = \frac{\epsilon_j}{k^2 - \chi_{ij} + i\epsilon} \quad (i \neq j). \quad (2)$$

The factor χ_{ij} is defined in terms of the squared masses as $\chi_{ij} = (\chi_i + \chi_j)/2$. The sign vector ϵ_j is defined as $+1$ for the fermionic valence and sea quarks ($j = 1, \dots, 6$) and -1 for the bosonic ghost quarks ($j = 7, 8, 9$). The propagator of a flavor-neutral meson in supersymmetric PQ χ PT is more complicated, as it connects mesons of different flavor indices as well. The propagator for a meson with quantum numbers $q_i \bar{q}_i$ to one with quantum numbers $q_j \bar{q}_j$ may be written in Minkowski space [5] as

$$G_{ij}^n(k) = G_{ij}^c(k) \delta_{ij} - \frac{1}{n_{\text{sea}}} G_{ij}^q(k). \quad (3)$$

The nontrivial part G_{ij}^q of the neutral meson propagator may be expressed, by means of partial fractioning, in terms of a sum of single and double poles in k^2 . For the most general case of $d_{\text{sea}} = 3$, in terms of the single-pole residue R , the double-pole residue R^d and the auxiliary residue R^c , the propagator G_{ij}^q is

$$-i G_{ij}^q(k) = \frac{R_{j\pi\eta}^i}{k^2 - \chi_i + i\epsilon} + \frac{R_{i\pi\eta}^j}{k^2 - \chi_j + i\epsilon} \quad (4)$$

$$+ \frac{R_{\eta ij}^\pi}{k^2 - \chi_\pi + i\epsilon} + \frac{R_{\pi ij}^\eta}{k^2 - \chi_\eta + i\epsilon} \quad (i \neq j),$$

$$\begin{aligned} -i G_{ii}^q(k) &= \frac{R_i^d}{(k^2 - \chi_i + i\epsilon)^2} + \frac{R_i^c}{k^2 - \chi_i + i\epsilon} \\ &+ \frac{R_{\eta ii}^\pi}{k^2 - \chi_\pi + i\epsilon} + \frac{R_{\pi ii}^\eta}{k^2 - \chi_\eta + i\epsilon}. \end{aligned} \quad (5)$$

For uniformity of notation, the squared (lowest order) masses of the neutral pion and the eta meson in the sea quark sector have been denoted by χ_π and χ_η , respectively. They are functions of the sea quark masses, and in the case of $d_{\text{sea}} = 3$ they are given by the lowest order χ PT result with $\pi^0 - \eta$ mixing active [5]. They can be obtained from the relations

$$\begin{aligned} \chi_\pi + \chi_\eta &= \frac{2}{3} (\chi_4 + \chi_5 + \chi_6), \\ \chi_\pi \chi_\eta &= \frac{1}{3} (\chi_4 \chi_5 + \chi_5 \chi_6 + \chi_4 \chi_6). \end{aligned} \quad (6)$$

For $d_{\text{sea}} = 2$, the π^0 pole in G_{ij}^q disappears, and in that case the (lowest order) eta meson mass is trivially given in terms of the remaining sea quark masses as $\chi_\eta = 1/3 \chi_4 + 2/3 \chi_6$. In this case the index π has been suppressed in the residue notation. Further, for $d_{\text{sea}} = 1$, both the π^0 and η poles disappear and consequently both the π and η indices have been suppressed. In this way we obtain a unique notation for the residues, from which the number of nondegenerate sea quarks is immediately apparent.

The various residues R of the propagators of the neutral, or flavor-diagonal mesons [4, 5] appear in the results, and are one reason why the PQ χ PT expressions are much more involved than the χ PT results of Ref. [13]. The use of the propagators (4) and (5) in the present form has the advantage of producing results in terms of standard loop integrals which can be treated with known methods. On the other hand, the various residues R of the flavor-neutral meson propagator fulfill a very large number of relations and the direct output from the calculations of the diagrams consequently produces a large number of redundant terms. This problem does not yet manifest itself at the one-loop level, but becomes troublesome at the two-loop level when the mass degeneracies in the sea and valence quark sectors are lifted. With a major effort, the end result can be simplified, in some cases it has been compressed by more than an order of magnitude.

C. Notation for Propagator Residues

The form of the single-pole residues R_{jkl}^i and the double-pole residue R_i^d , which appear in Eqs. (4) and (5), depends on the degree of degeneracy in the sea quark masses, which in turn is indicated by the number of indices in the single-pole residue R . It is useful to define the more general quantities $R_{a\dots b}^z$ such that

$$\begin{aligned} R_{ab}^z &= \chi_a - \chi_b, \\ R_{abc}^z &= \frac{\chi_a - \chi_b}{\chi_a - \chi_c}, \\ R_{abcd}^z &= \frac{(\chi_a - \chi_b)(\chi_a - \chi_c)}{\chi_a - \chi_d}, \\ R_{abcdefg}^z &= \frac{(\chi_a - \chi_b)(\chi_a - \chi_c)(\chi_a - \chi_d)}{(\chi_a - \chi_e)(\chi_a - \chi_f)(\chi_a - \chi_g)}, \end{aligned} \quad (7)$$

and so on. The R^z notation is primarily useful for defining the residues of the flavor-neutral propagators, but it may also appear independently in the final result for the decay constant. In such cases, the R^z have been generated by simplification procedures, as all the residues that are naturally generated by partial fractioning of the propagator G_{ij}^q are of the form given below. Note that $R_{a\dots b}^z$ has the same dimension as χ_i for an even number of indices and is dimensionless for an odd number of indices.

For the case of $d_{\text{sea}} = 1$, all residues associated with the sea quark sector have reduced to numbers. Some

nontrivial residues still appear if the valence quarks are nondegenerate, according to

$$\begin{aligned} R_j^i &= R_{i4j}^z, \\ R_i^d &= R_{i4}^z. \end{aligned} \quad (8)$$

For convenience, R_i^d is also used for $d_{\text{val}} = 1$ in order to maintain a notation similar to the more general cases.

As noted in Ref. [5], the residues also simplify for $d_{\text{sea}} = 2$. In particular, the residue of the neutral pion pole in the sea quark sector (R_{jkl}^π) vanishes, and the remaining ones satisfy a larger number of relations than for $d_{\text{sea}} = 3$. The remaining nontrivial residues for $d_{\text{sea}} = 2$ may be expressed as

$$\begin{aligned} R_{jk}^i &= R_{i46jk}^z, \\ R_i^d &= R_{i46\eta}^z, \\ R_i^c &= R_{4\eta}^i + R_{6\eta}^i - R_{\eta\eta}^i. \end{aligned} \quad (9)$$

It is also noteworthy that the above residue notation is highly redundant because of the trivial relation between χ_4, χ_6 and χ_η . This fact has been exploited in the simplification of the end results.

For $d_{\text{sea}} = 3$, the naturally generated residues are

$$\begin{aligned} R_{jkl}^i &= R_{i456jkl}^z, \\ R_i^d &= R_{i456\pi\eta}^z, \\ R_i^c &= R_{4\pi\eta}^i + R_{5\pi\eta}^i + R_{6\pi\eta}^i - R_{\pi\eta\eta}^i - R_{\pi\pi\eta}^i. \end{aligned} \quad (10)$$

Even in this case there are many relations between the various residues. Some can be found in Ref. [4] but many more exist. Finally, it should be noted that in the limit where the sea quark masses become equal to the valence quark masses, the propagator residues of PQ χ PT reduce so that the π^0 and η meson propagators of unquenched χ PT are recovered.

D. Integral Notation

The expression for the decay constant of a charged pseudoscalar meson depends, in NNLO PQ χ PT, on a number of one-loop and two-loop integrals. The finite parts of the chiral logarithms A, B and C are

$$\begin{aligned} \bar{A}(\chi) &= -\pi_{16} \chi \log(\chi/\mu^2), \\ \bar{B}(\chi_i, \chi_j; 0) &= -\pi_{16} \frac{\chi_i \log(\chi_i/\mu^2) - \chi_j \log(\chi_j/\mu^2)}{\chi_i - \chi_j}, \\ \bar{C}(\chi, \chi, \chi; 0) &= -\pi_{16}/(2\chi), \end{aligned} \quad (11)$$

where the subtraction scale dependence has been moved into the loop integrals. We also define $\pi_{16} = 1/(16\pi^2)$. Note that in the limit $\chi_i = \chi_j$, the integral $\bar{B}$ reduces to

$$\bar{B}(\chi, \chi; 0) = -\pi_{16} (1 + \log(\chi/\mu^2)). \quad (12)$$

The following combinations of finite one-loop integrals are also introduced, as they are naturally generated by the procedure of dimensional regularization:

$$\begin{aligned}\bar{A}(\chi; \varepsilon) &= \bar{A}(\chi)^2 / (2\pi_{16} \chi) \\ &+ \pi_{16} \chi (\pi^2/12 + 1/2), \\ \bar{B}(\chi, \chi; 0, \varepsilon) &= \bar{A}(\chi) \bar{B}(\chi, \chi; 0) / (\pi_{16} \chi) \\ &- \bar{A}(\chi)^2 / (2\pi_{16} \chi^2) \\ &+ \pi_{16} (\pi^2/12 + 1/2).\end{aligned}\quad (13)$$

Additionally, the upper middle and upper left diagrams in Fig. 1 generate integrals of the form

$$\bar{B}(\chi_i, \chi_j; 0, k) = \chi_i \bar{B}(\chi_i, \chi_j; 0) + \bar{A}(\chi_j), \quad (14)$$

which are symmetric under the interchange of χ_i and χ_j . The above integrals have been introduced in order to make the symmetries in the end results more explicit.

The finite two-loop integrals H^F, H_1^F, H_{21}^F that are generated by the top right diagram of Fig. 1 may be evaluated using the methods of Ref. [13]. Note that the corresponding primed integrals $H^{F'}, H_1^{F'}, H_{21}^{F'}$ indicate differentiation with respect to p^2 . The notation used for the H integrals in this paper is similar to that of Ref. [13], except that an extra integer argument now indicates the propagator structure, such that e.g.

$$H^F(\chi_i, \chi_j, \chi_k; p^2) \rightarrow H^F(n, \chi_i, \chi_j, \chi_k; p^2). \quad (15)$$

The case of $n = 1$ indicates that the integral consists of single propagators only, as in Ref. [13], whereas $n = 2$ indicates that the first propagator appears squared and $n = 3$ that the second propagator appears squared. The cases with two double propagators that can appear in the calculations are $n = 5$, for which the first and second propagators appear squared, and $n = 7$ for which the second and third propagators are squared. Explicit expressions for $n = 1$ can be found in Ref. [13], and the other cases may be obtained by differentiation with respect to the masses of those expressions.

E. Simplification of End Results

A number of combinations of quark masses and propagator residues are naturally generated in the calculations, and have consequently been given special notations. The most common of these is

$$\bar{\chi}_n = \frac{1}{3} \sum_{i=4,5,6} \chi_i^n, \quad (16)$$

of which $\bar{\chi}_1$ is equal to the average sea quark mass $\bar{\chi}$ defined in Ref. [4]. Other combinations that appear in the calculations consist of products of quark masses and propagator residues. For $d_{\text{sea}} = 2$, these include

$$\bar{\chi}_{bn}^a = \frac{1}{3} \sum_{i=4,5,6} R_{bi}^a \chi_i^n,$$

$$\begin{aligned}\bar{\chi}_{bdn}^{ac} &= \frac{1}{3} \sum_{i=4,5,6} R_{bi}^a R_{di}^c \chi_i^n, \\ \bar{\chi}_{bdn}^{ace} &= \frac{1}{3} \sum_{i=4,5,6} R_{bi}^a R_{di}^c R_{ii}^e \chi_i^n, \\ R_{ijk}^v &= R_{jj}^i + R_{kk}^i - 2R_{jk}^i.\end{aligned}\quad (17)$$

For $d_{\text{sea}} = 3$ these become

$$\begin{aligned}\bar{\chi}_{bn}^a &= \frac{1}{3} \sum_{i=4,5,6} R_{bii}^a \chi_i^n, \\ \bar{\chi}_{bcn}^a &= \frac{1}{3} \sum_{i=4,5,6} R_{bci}^a \chi_i^n, \\ \bar{\chi}_{bcefn}^{ad} &= \frac{1}{3} \sum_{i=4,5,6} R_{bci}^a R_{efi}^d \chi_i^n, \\ \bar{\chi}_{bcefn}^{adk} &= \frac{1}{3} \sum_{i=4,5,6} R_{bci}^a R_{efi}^d R_{lii}^k \chi_i^n, \\ R_{ijk}^v &= R_{jkk}^i + R_{jll}^i - 2R_{jkl}^i.\end{aligned}\quad (18)$$

It should be noted that also in the case of $d_{\text{sea}} = 2$, the sums run over all three sea quark flavors. Furthermore, only some of the above $\bar{\chi}$ functions appear in the results, as many of them can be reexpressed in terms of other R or R^z functions.

The actual simplification of the end results is at first glance a formidable task, since the expression for F_a as calculated from the diagrams in Fig. 1 contains several thousand terms. Especially for $d_{\text{val}} = 2$, the parts proportional to $\bar{A}(\chi_\eta; \varepsilon)$ and $\bar{A}(\chi_1; \varepsilon)$ alone are several hundred terms long, while in simplified form they typically contain only ~ 5 terms. However, these simplifications are not easily apparent and are consequently best accomplished by the employment of suitable software, such as **Maple** or **Mathematica**. Even so, as the expressions can seldom be factored into a single term, considerable trial and error is usually required.

F. Summation Conventions

The expressions for the decay constants are symmetric, within the sea and valence quark sectors, under the interchange of quark masses. Consequently, the end results can be conveniently compactified by the introduction of summation conventions which exploit these symmetries. For example, the notation for the sea quark sector may be considerably compactified by the introduction of the summation indices s and t , which may appear for the quark masses χ , and among the indices of the functions R and $\bar{\chi}$.

These sea-quark summation indices should be interpreted as follows: If an index t is present once or several times, then there will always be an occurrence of the s index as well, and the term is then to be summed over all pairs of different sea quark indices. If the index s is

present but t is not, then the entire term is to be summed over all sea quark indices. Simple examples of terms are

$$\begin{aligned}\chi_s &= \sum_{i=4,5,6} \chi_i, \\ \chi_{st} &= \sum_{\substack{i,j=4,5,6 \\ j>i}} \chi_{ij}.\end{aligned}\quad (19)$$

For terms consisting of products of several factors, the summation sign should always be inserted at the beginning of each relevant term, such that e.g.

$$\bar{A}(\chi_s) R_{\eta s}^1 \chi_s = \sum_{i=4,5,6} \bar{A}(\chi_i) R_{\eta i}^1 \chi_i. \quad (20)$$

Thus any contribution which is written in terms of the indices s and t explicitly fulfills the required symmetry properties in the sea quark sector. Note that also for $d_{\text{sea}} = 2$, the summation is over all three sea quark flavors, although the index t has not been implemented in that case.

Further compactification of the results is possible, as the valence quark sector is symmetric under the exchange of the valence quark indices 1 and 3. For this purpose, the summation indices p and q have been introduced. These are in the present paper only needed for the case of $d_{\text{val}} = 2$ and then always occur for the squared valence quark masses χ_1 and χ_3 . If the index q is present, there will always be an index p and the resulting sum is over the pairs $(p, q) = (1, 3)$ and $(p, q) = (3, 1)$. If only p is present, the sum is over the indices 1 and 3. An example is

$$\bar{A}(\chi_p) R_{q\eta}^p \chi_p = \bar{A}(\chi_1) R_{3\eta}^1 \chi_1 + [1 \leftrightarrow 3]. \quad (21)$$

Any contribution written in terms of the (p, q) notation is thus symmetric under the interchange of the valence quark masses χ_1 and χ_3 .

For $d_{\text{sea}} = 3$, there exists an additional symmetry, i.e. the results are symmetric under the interchange of the lowest order neutral meson masses, χ_π and χ_η , in the sea-quark sector. This is exploited by the summation indices m and n . If the index m is present, there will always be an index n and the corresponding sum is over the pairs $(m, n) = (\pi, \eta)$ and $(m, n) = (\eta, \pi)$. An example is

$$\bar{A}(\chi_m) R_{n11}^m \chi_m = \bar{A}(\chi_\eta) R_{\pi11}^\eta \chi_\eta + [\eta \leftrightarrow \pi]. \quad (22)$$

The above summation conventions thus provide a means of eliminating a large number of terms in the expression for F_a , which are of similar form. As an added bonus, it gives a possibility to conveniently check that the NNLO loop results fulfill the required symmetry relations in the sea and valence sectors.

III. THE PSEUDOSCALAR MESON DECAY CONSTANTS TO $\mathcal{O}(p^6)$

The decay constants F_a of the pseudoscalar mesons are obtained from the definition

$$\langle 0 | A_a^\mu(0) | \phi_a(p) \rangle = i\sqrt{2} p^\mu F_a, \quad (23)$$

in terms of the axial current operator A_a^μ . The diagrams that contribute to that operator at $\mathcal{O}(p^6)$, or NNLO, are shown in Fig. 1. Diagrams at $\mathcal{O}(p^2)$ and $\mathcal{O}(p^4)$ also contribute to Eq. (23) via the renormalization of the pseudoscalar meson wave function $\phi_a(p)$. The results for charged pseudoscalar mesons so obtained depend on the $\mathcal{O}(p^6)$ low-energy constants K_{19}^r through K_{23}^r .




FIG. 1: Feynman diagrams at $\mathcal{O}(p^6)$ or two-loop for the matrix element of the axial current operator. Filled circles denote vertices of the $\mathcal{L}_2$ Lagrangian, whereas open squares and shaded diamonds denote vertices of the $\mathcal{L}_4$ and $\mathcal{L}_6$ Lagrangians, respectively.

The results are expressed in terms of the valence inputs χ_1, χ_3 and the sea inputs χ_4, χ_5, χ_6 , which are defined in terms of the quark masses through $\chi_i = 2B_0 m_i$, and the quantity $\chi_{ij} = (\chi_i + \chi_j)/2$, which corresponds to the lowest order charged meson mass. Other parameters include the decay constant in the chiral limit (F_0), the quark condensate in the chiral limit, via $\langle \bar{q}q \rangle = -B_0 F_0^2$, and the LEC:s of $\mathcal{O}(p^4)$ and $\mathcal{O}(p^6)$, i.e. the L_i^r and the K_i^r [11, 12], respectively.

The decay constants of the pseudoscalar mesons are given in the form

$$F_{\text{phys}} = F_0 \left[1 + \frac{\delta^{(4)\text{vs}}}{F_0^2} + \frac{\delta_{\text{ct}}^{(6)\text{vs}} + \delta_{\text{loops}}^{(6)\text{vs}}}{F_0^4} + \mathcal{O}(p^8) \right], \quad (24)$$

where the $\mathcal{O}(p^4)$ and $\mathcal{O}(p^6)$ contributions have been separated. The NNLO contribution $\delta^{(6)}$ has been further split into the contributions from the chiral loops and from the $\mathcal{O}(p^6)$ counterterms. The superscripts (v) and (s) indicate the values of d_{val} and d_{sea} , respectively.

A. Results for $d_{\text{val}} = 1$

The NLO result for $d_{\text{val}} = 1$ is fairly short, and will thus only be given for $d_{\text{sea}} = 3$. The results for $d_{\text{sea}} = 1, 2$ may readily be derived from that expression. The combined NLO result (loops and counterterms) for $d_{\text{sea}} = 3$, which is in agreement with Ref. [5], is

$$\delta^{(4)13} = 12 L_4^r \bar{\chi}_1 + 4 L_5^r \chi_1 + 1/2 \bar{A}(\chi_{1s}). \quad (25)$$

At NNLO, the chiral loops form, by far, the largest contribution to the decay constant. As a straightforward derivation of the results for $d_{\text{sea}} = 1, 2$ from the $d_{\text{sea}} = 3$ case is tedious and complicated, the expressions for the different cases are given separately below. As expected, the infinities in all expressions for the decay constant have canceled. The appearance of 'unphysical' $\bar{B}$ logarithms

in the results is, in part, due to the partial quenching, and to the fact that the results have been expressed in terms of the lowest order masses rather than the full physical ones. Consequently, not all of them correspond to the quenched chiral logarithms which are ill-behaved in the chiral limit.

The contribution from the $\mathcal{O}(p^6)$ counterterms to the decay constant at NNLO is, for $d_{\text{sea}} = 3$,

$$\begin{aligned} \delta_{\text{ct}}^{(6)13} = & 8 K_{19}^r \chi_1^2 + 24 K_{20}^r \bar{\chi}_1 \chi_1 + 24 K_{21}^r \bar{\chi}_2 \\ & + 72 K_{22}^r \bar{\chi}_1^2 + 8 K_{23}^r \chi_1^2, \end{aligned} \quad (26)$$

from which the results for $d_{\text{sea}} = 1, 2$ may be readily inferred. The NNLO contributions from the chiral loops for $d_{\text{sea}} = 1, 2, 3$ are, respectively,

$$\begin{aligned} \delta_{\text{loops}}^{(6)11} = & \pi_{16} L_0^r [-13/3 \chi_1 \chi_4 + 1/2 \chi_1^2 - 3/2 \chi_4^2] - 2 \pi_{16} L_1^r \chi_1^2 - \pi_{16} L_2^r [\chi_1^2 + 8 \chi_4^2] \\ & + \pi_{16} L_3^r [-17/6 \chi_1 \chi_4 + 5/4 \chi_1^2 - 3/4 \chi_4^2] + \pi_{16}^2 [-1/2 \chi_1 \chi_4 + 1/64 \chi_1^2 - 73/128 \chi_4^2] \\ & - 48 L_4^r L_5^r \chi_1 \chi_4 - 72 L_4^{r2} \chi_4^2 - 8 L_5^{r2} \chi_1^2 + 4 \bar{A}(\chi_1) L_0^r [\chi_1 + R_1^d] - 4 \bar{A}(\chi_1) L_1^r \chi_1 - 10 \bar{A}(\chi_1) L_2^r \chi_1 \\ & + 4 \bar{A}(\chi_1) L_3^r [\chi_1 + R_1^d] - 4/3 \bar{A}(\chi_1) L_5^r \chi_1 - 1/2 \bar{A}(\chi_1) \bar{B}(\chi_{14}, \chi_{14}; 0) \chi_{14} + 1/8 \bar{A}(\chi_1; \varepsilon) \pi_{16} \chi_4 \\ & - \bar{A}(\chi_{14}) \pi_{16} [3/4 \chi_{14} + 9/8 \chi_4] - 12 \bar{A}(\chi_{14}) L_0^r \chi_{14} - 30 \bar{A}(\chi_{14}) L_3^r \chi_{14} - 18 \bar{A}(\chi_{14}) L_4^r \chi_4 \\ & + 6 \bar{A}(\chi_{14}) L_5^r \chi_1 + 9/4 \bar{A}(\chi_{14}; \varepsilon) \pi_{16} \chi_4 - 64 \bar{A}(\chi_4) L_1^r \chi_4 - 16 \bar{A}(\chi_4) L_2^r \chi_4 + 32 \bar{A}(\chi_4) L_4^r \chi_4 \\ & + \bar{A}(\chi_4; \varepsilon) \pi_{16} \chi_4 + 4 \bar{B}(\chi_1, \chi_1; 0) L_0^r R_1^d \chi_1 + 4 \bar{B}(\chi_1, \chi_1; 0) L_3^r R_1^d \chi_1 - 4/3 \bar{B}(\chi_1, \chi_1; 0) L_5^r R_1^d \chi_1 \\ & - 1/8 \bar{B}(\chi_1, \chi_1; 0, \varepsilon) \pi_{16} R_1^d \chi_1 - 36 \bar{B}(\chi_{14}, \chi_{14}; 0) L_4^r \chi_{14} \chi_4 - 12 \bar{B}(\chi_{14}, \chi_{14}; 0) L_5^r \chi_{14}^2 \\ & + 72 \bar{B}(\chi_{14}, \chi_{14}; 0) L_6^r \chi_{14} \chi_4 + 24 \bar{B}(\chi_{14}, \chi_{14}; 0) L_8^r \chi_{14}^2 - 1/8 H^F(1, \chi_1, \chi_{14}, \chi_{14}; \chi_1) [\chi_1 - R_1^d] \\ & - H^F(1, \chi_{14}, \chi_{14}, \chi_4; \chi_1) \chi_4 + 1/8 H^F(2, \chi_1, \chi_{14}, \chi_{14}; \chi_1) R_1^d \chi_1 + 5/18 H^{F'}(1, \chi_1, \chi_1, \chi_1; \chi_1) \chi_1^2 \\ & + H^{F'}(1, \chi_1, \chi_{14}, \chi_{14}; \chi_1) [1/8 \chi_1 \chi_4 - 1/2 \chi_1^2] + H^{F'}(1, \chi_{14}, \chi_{14}, \chi_4; \chi_1) \chi_1 \chi_4 \\ & + 2/9 H^{F'}(2, \chi_1, \chi_1, \chi_1; \chi_1) R_1^d \chi_1^2 + 3/8 H^{F'}(2, \chi_1, \chi_{14}, \chi_{14}; \chi_1) R_1^d \chi_1^2 + 1/9 H^{F'}(5, \chi_1, \chi_1, \chi_1; \chi_1) (R_1^d)^2 \chi_1^2 \\ & - 2 H_1^{F'}(3, \chi_{14}, \chi_1, \chi_{14}; \chi_1) R_1^d \chi_1^2 + 3/8 H_{21}^{F'}(1, \chi_1, \chi_{14}, \chi_{14}; \chi_1) \chi_1^2 + 3 H_{21}^{F'}(1, \chi_4, \chi_{14}, \chi_{14}; \chi_1) \chi_1^2 \\ & - 3/8 H_{21}^{F'}(2, \chi_1, \chi_{14}, \chi_{14}; \chi_1) R_1^d \chi_1^2, \end{aligned} \quad (27)$$

$$\begin{aligned} \delta_{\text{loops}}^{(6)12} = & \pi_{16} L_0^r [4/9 \chi_\eta \chi_4 + 1/2 \chi_1^2 - 13/3 \bar{\chi}_1 \chi_1 - 35/18 \bar{\chi}_2] - 2 \pi_{16} L_1^r \chi_1^2 \\ & - \pi_{16} L_2^r [11/3 \chi_\eta \chi_4 + \chi_1^2 + 13/3 \bar{\chi}_2] + \pi_{16} L_3^r [4/9 \chi_\eta \chi_4 + 5/4 \chi_1^2 - 17/6 \bar{\chi}_1 \chi_1 - 43/36 \bar{\chi}_2] \\ & + \pi_{16}^2 [-15/64 \chi_\eta \chi_4 + 1/64 \chi_1^2 - 1/2 \bar{\chi}_1 \chi_1 - 43/128 \bar{\chi}_2] - 48 L_4^r L_5^r \bar{\chi}_1 \chi_1 - 72 L_4^{r2} \bar{\chi}_1^2 - 8 L_5^{r2} \chi_1^2 \\ & + 4 \bar{A}(\chi_\eta) L_0^r R_{11}^\eta \chi_\eta - 8 \bar{A}(\chi_\eta) L_1^r \chi_\eta - 2 \bar{A}(\chi_\eta) L_2^r \chi_\eta + 4 \bar{A}(\chi_\eta) L_3^r R_{11}^\eta \chi_\eta + 4 \bar{A}(\chi_\eta) L_4^r \chi_\eta \\ & - 4/3 \bar{A}(\chi_\eta) L_5^r R_{11}^\eta \chi_1 - 1/6 \bar{A}(\chi_\eta) \bar{B}(\chi_{1s}, \chi_{1s}; 0) R_{1s}^\eta \chi_{1s} + 1/8 \bar{A}(\chi_\eta; \varepsilon) \pi_{16} R_{1s}^c \chi_\eta \\ & + 4 \bar{A}(\chi_1) L_0^r [R_1^c \chi_1 + R_1^d] - 4 \bar{A}(\chi_1) L_1^r \chi_1 - 10 \bar{A}(\chi_1) L_2^r \chi_1 + 4 \bar{A}(\chi_1) L_3^r [R_1^c \chi_1 + R_1^d] \\ & - 4/3 \bar{A}(\chi_1) L_5^r R_{1s}^c \chi_1 - 1/6 \bar{A}(\chi_1) \bar{B}(\chi_{1s}, \chi_{1s}; 0) R_{s\eta}^1 \chi_{1s} + \bar{A}(\chi_1; \varepsilon) \pi_{16} [1/4 \chi_1 - 1/8 R_{1s}^c \chi_1 - 1/8 R_1^d] \\ & - 24 \bar{A}(\chi_4) L_1^r \chi_4 - 6 \bar{A}(\chi_4) L_2^r \chi_4 + 12 \bar{A}(\chi_4) L_4^r \chi_4 + 3/8 \bar{A}(\chi_4; \varepsilon) \pi_{16} \chi_4 - 32 \bar{A}(\chi_{46}) L_1^r \chi_{46} \\ & - 8 \bar{A}(\chi_{46}) L_2^r \chi_{46} + 16 \bar{A}(\chi_{46}) L_4^r \chi_{46} + 1/2 \bar{A}(\chi_{46}; \varepsilon) \pi_{16} \chi_{46} - \bar{A}(\chi_{1s}) \pi_{16} [1/4 \chi_{1s} + 3/8 \bar{\chi}_1] \\ & - 4 \bar{A}(\chi_{1s}) L_0^r \chi_{1s} - 10 \bar{A}(\chi_{1s}) L_3^r \chi_{1s} - 6 \bar{A}(\chi_{1s}) L_4^r \bar{\chi}_1 + 2 \bar{A}(\chi_{1s}) L_5^r \chi_1 + 3/8 \bar{A}(\chi_{1s}; \varepsilon) \pi_{16} [\chi_s + \bar{\chi}_1] \\ & + 4 \bar{B}(\chi_1, \chi_1; 0) L_0^r R_1^d \chi_1 + 4 \bar{B}(\chi_1, \chi_1; 0) L_3^r R_1^d \chi_1 - 4/3 \bar{B}(\chi_1, \chi_1; 0) L_5^r R_1^d \chi_1 \\ & - 1/8 \bar{B}(\chi_1, \chi_1; 0, \varepsilon) \pi_{16} R_1^d \chi_1 - 12 \bar{B}(\chi_{1s}, \chi_{1s}; 0) L_4^r \bar{\chi}_1 \chi_{1s} - 4 \bar{B}(\chi_{1s}, \chi_{1s}; 0) L_5^r \chi_{1s}^2 \\ & + 24 \bar{B}(\chi_{1s}, \chi_{1s}; 0) L_6^r \bar{\chi}_1 \chi_{1s} + 8 \bar{B}(\chi_{1s}, \chi_{1s}; 0) L_8^r \chi_{1s}^2 + 1/24 H^F(1, \chi_\eta, \chi_{1s}, \chi_{1s}; \chi_1) R_{\eta 1s}^v \chi_\eta \end{aligned}$$

$$\begin{aligned}
& + H^F(1, \chi_1, \chi_{1s}, \chi_{1s}; \chi_1) [-1/12 R_{s\eta}^1 \chi_1 + 1/24 R_1^c \chi_1 + 1/24 R_1^d] - 3/8 H^F(1, \chi_{14}, \chi_{14}, \chi_4; \chi_1) \chi_4 \\
& - 1/2 H^F(1, \chi_{14}, \chi_{16}, \chi_{46}; \chi_1) \chi_{46} + 1/24 H^F(2, \chi_1, \chi_{1s}, \chi_{1s}; \chi_1) R_1^d \chi_1 \\
& + 1/9 H^{F'}(1, \chi_\eta, \chi_\eta, \chi_1; \chi_1) (R_{11}^\eta)^2 \chi_1^2 + 2/9 H^{F'}(1, \chi_\eta, \chi_1, \chi_1; \chi_1) R_{11}^\eta R_1^c \chi_1^2 \\
& - H^{F'}(1, \chi_\eta, \chi_{1s}, \chi_{1s}; \chi_1) [1/6 R_{11}^\eta \chi_1^2 + 1/24 R_{\eta 1s}^v \chi_\eta \chi_1] + H^{F'}(1, \chi_1, \chi_1, \chi_1; \chi_1) [1/6 \chi_1^2 \\
& + 1/9 (R_1^c)^2 \chi_1^2] + H^{F'}(1, \chi_1, \chi_{1s}, \chi_{1s}; \chi_1) [-1/4 R_{s\eta}^1 \chi_1^2 + 1/8 R_1^c \chi_1^2 - 1/24 R_1^d \chi_1] \\
& + 3/8 H^{F'}(1, \chi_{14}, \chi_{14}, \chi_4; \chi_1) \chi_1 \chi_4 + 1/2 H^{F'}(1, \chi_{14}, \chi_{16}, \chi_{46}; \chi_1) \chi_1 \chi_{46} \\
& + 2/9 H^{F'}(2, \chi_1, \chi_\eta, \chi_1; \chi_1) R_{11}^\eta R_1^d \chi_1^2 + 2/9 H^{F'}(2, \chi_1, \chi_1, \chi_1; \chi_1) R_1^c R_1^d \chi_1^2 \\
& + 1/8 H^{F'}(2, \chi_1, \chi_{1s}, \chi_{1s}; \chi_1) R_1^d \chi_1^2 + 1/9 H^{F'}(5, \chi_1, \chi_1, \chi_1; \chi_1) (R_1^d)^2 \chi_1^2 \\
& - 1/3 H_1^{F'}(1, \chi_\eta, \chi_{1s}, \chi_{1s}; \chi_1) R_{1s}^\eta R_{1s\eta}^z \chi_1^2 - 2/3 H_1^{F'}(1, \chi_{1s}, \chi_{1s}, \chi_1; \chi_1) R_{1s}^\eta R_{1s\eta}^z \chi_1^2 \\
& - 2/3 H_1^{F'}(3, \chi_{1s}, \chi_1, \chi_{1s}; \chi_1) R_1^d \chi_1^2 - 1/8 H_{21}^{F'}(1, \chi_\eta, \chi_{1s}, \chi_{1s}; \chi_1) R_{\eta 1s}^v \chi_1^2 \\
& + H_{21}^{F'}(1, \chi_1, \chi_{1s}, \chi_{1s}; \chi_1) [1/4 R_{s\eta}^1 \chi_1^2 - 1/8 R_1^c \chi_1^2] + 9/8 H_{21}^{F'}(1, \chi_4, \chi_{14}, \chi_{14}; \chi_1) \chi_1^2 \\
& + 3/2 H_{21}^{F'}(1, \chi_{46}, \chi_{14}, \chi_{16}; \chi_1) \chi_1^2 - 1/8 H_{21}^{F'}(2, \chi_1, \chi_{1s}, \chi_{1s}; \chi_1) R_1^d \chi_1^2, \tag{28}
\end{aligned}$$

$$\begin{aligned}
\delta_{\text{loops}}^{(6)13} = & \pi_{16} L_0^r [4/9 \chi_\pi \chi_\eta + 1/2 \chi_1^2 - 13/3 \bar{\chi}_1 \chi_1 - 35/18 \bar{\chi}_2] - 2 \pi_{16} L_1^r \chi_1^2 \\
& - \pi_{16} L_2^r [11/3 \chi_\pi \chi_\eta + \chi_1^2 + 13/3 \bar{\chi}_2] + \pi_{16} L_3^r [4/9 \chi_\pi \chi_\eta + 5/4 \chi_1^2 - 17/6 \bar{\chi}_1 \chi_1 - 43/36 \bar{\chi}_2] \\
& + \pi_{16}^2 [-15/64 \chi_\pi \chi_\eta + 1/64 \chi_1^2 - 1/2 \bar{\chi}_1 \chi_1 - 43/128 \bar{\chi}_2] - 48 L_4^r L_5^r \bar{\chi}_1 \chi_1 - 72 L_4^r \bar{\chi}_1^2 - 8 L_5^r \chi_1^2 \\
& + 4 \bar{A}(\chi_m) L_0^r R_{n11}^m \chi_m + 8 \bar{A}(\chi_m) L_1^r \bar{\chi}_{n0}^m \chi_m + 2 \bar{A}(\chi_m) L_2^r \bar{\chi}_{n0}^m \chi_m + 4 \bar{A}(\chi_m) L_3^r R_{n11}^m \chi_m \\
& - 4 \bar{A}(\chi_m) L_4^r \bar{\chi}_{n1}^m - 4/3 \bar{A}(\chi_m) L_5^r R_{n11}^m \chi_1 - 1/6 \bar{A}(\chi_m) \bar{B}(\chi_{1s}, \chi_{1s}, 0) R_{n1s}^m \chi_{1s} \\
& + \bar{A}(\chi_m; \varepsilon) \pi_{16} [-7/24 \bar{\chi}_{n0}^m \chi_m + 1/6 \bar{\chi}_{n1}^m - 1/8 R_{n11}^m \chi_m] + 4 \bar{A}(\chi_1) L_0^r [R_1^c \chi_1 + R_1^d] \\
& - 4 \bar{A}(\chi_1) L_1^r \chi_1 - 10 \bar{A}(\chi_1) L_2^r \chi_1 + 4 \bar{A}(\chi_1) L_3^r [R_1^c \chi_1 + R_1^d] - 4/3 \bar{A}(\chi_1) L_5^r R_1^c \chi_1 \\
& - 1/6 \bar{A}(\chi_1) \bar{B}(\chi_{1s}, \chi_{1s}, 0) R_{s\pi\eta}^1 \chi_{1s} + \bar{A}(\chi_1; \varepsilon) \pi_{16} [1/4 \chi_1 - 1/8 R_1^c \chi_1 - 1/8 R_1^d] \\
& - \bar{A}(\chi_{1s}) \pi_{16} [1/4 \chi_{1s} + 3/8 \bar{\chi}_1] - 4 \bar{A}(\chi_{1s}) L_0^r \chi_{1s} - 10 \bar{A}(\chi_{1s}) L_3^r \chi_{1s} - 6 \bar{A}(\chi_{1s}) L_4^r \bar{\chi}_1 \\
& + 2 \bar{A}(\chi_{1s}) L_5^r \chi_1 + 3/8 \bar{A}(\chi_{1s}; \varepsilon) \pi_{16} [\chi_s + \bar{\chi}_1] + \bar{A}(\chi_s) L_1^r [-8 \chi_s + 8/3 R_s^c \chi_s] \\
& - \bar{A}(\chi_s) L_2^r [2 \chi_s - 2/3 R_s^c \chi_s] + \bar{A}(\chi_s) L_4^r [4 \chi_s - 4/3 R_s^c \chi_s] + \bar{A}(\chi_s; \varepsilon) \pi_{16} [1/8 \chi_s - 1/24 R_s^c \chi_s] \\
& - 16 \bar{A}(\chi_{st}) L_1^r \chi_{st} - 4 \bar{A}(\chi_{st}) L_2^r \chi_{st} + 8 \bar{A}(\chi_{st}) L_4^r \chi_{st} + 1/4 \bar{A}(\chi_{st}; \varepsilon) \pi_{16} \chi_{st} \\
& + 4 \bar{B}(\chi_1, \chi_1; 0) L_0^r R_1^d \chi_1 + 4 \bar{B}(\chi_1, \chi_1; 0) L_3^r R_1^d \chi_1 - 4/3 \bar{B}(\chi_1, \chi_1; 0) L_5^r R_1^d \chi_1 \\
& - 1/8 \bar{B}(\chi_1, \chi_1; 0, \varepsilon) \pi_{16} R_1^d \chi_1 - 12 \bar{B}(\chi_{1s}, \chi_{1s}; 0) L_4^r \bar{\chi}_1 \chi_{1s} - 4 \bar{B}(\chi_{1s}, \chi_{1s}; 0) L_5^r \chi_{1s}^2 \\
& + 24 \bar{B}(\chi_{1s}, \chi_{1s}; 0) L_6^r \bar{\chi}_1 \chi_{1s} + 8 \bar{B}(\chi_{1s}, \chi_{1s}; 0) L_8^r \chi_{1s}^2 + 1/24 H^F(1, \chi_m, \chi_{1s}, \chi_{1s}; \chi_1) R_{mn1s}^v \chi_m \\
& + H^F(1, \chi_1, \chi_{1s}, \chi_{1s}; \chi_1) [-1/12 R_{s\pi\eta}^1 \chi_1 + 1/24 R_1^c \chi_1 + 1/24 R_1^d] + H^F(1, \chi_{1s}, \chi_{1s}, \chi_s; \chi_1) [-1/8 \chi_s \\
& + 1/24 R_s^c \chi_s] - 1/4 H^F(1, \chi_{1s}, \chi_{1t}, \chi_{st}; \chi_1) \chi_{st} + 1/24 H^F(2, \chi_1, \chi_{1s}, \chi_{1s}; \chi_1) R_1^d \chi_1 \\
& + 1/9 H^{F'}(1, \chi_m, \chi_m, \chi_1; \chi_1) (R_{n11}^m)^2 \chi_1^2 + 2/9 H^{F'}(1, \chi_m, \chi_1, \chi_1; \chi_1) R_{n11}^m R_1^c \chi_1^2 \\
& - H^{F'}(1, \chi_m, \chi_{1s}, \chi_{1s}; \chi_1) [1/6 R_{n11}^m \chi_1^2 + 1/24 R_{mn1s}^v \chi_m \chi_1] \\
& + 2/9 H^{F'}(1, \chi_\pi, \chi_\eta, \chi_1; \chi_1) R_{\eta 11}^\pi R_{\pi 11}^\eta \chi_1^2 + H^{F'}(1, \chi_1, \chi_1, \chi_1; \chi_1) [1/6 \chi_1^2 + 1/9 (R_1^c)^2 \chi_1^2] \\
& + H^{F'}(1, \chi_1, \chi_{1s}, \chi_{1s}; \chi_1) [-1/4 R_{s\pi\eta}^1 \chi_1^2 + 1/8 R_1^c \chi_1^2 - 1/24 R_1^d \chi_1] + H^{F'}(1, \chi_{1s}, \chi_{1s}, \chi_s; \chi_1) [1/8 \chi_1 \chi_s \\
& - 1/24 R_s^c \chi_1 \chi_s] + 1/4 H^{F'}(1, \chi_{1s}, \chi_{1t}, \chi_{st}; \chi_1) \chi_1 \chi_{st} + 2/9 H^{F'}(2, \chi_1, \chi_m, \chi_1; \chi_1) R_{n11}^m R_1^d \chi_1^2 \\
& + 2/9 H^{F'}(2, \chi_1, \chi_1, \chi_1; \chi_1) R_1^c R_1^d \chi_1^2 + 1/8 H^{F'}(2, \chi_1, \chi_{1s}, \chi_{1s}; \chi_1) R_1^d \chi_1^2 \\
& + 1/9 H^{F'}(5, \chi_1, \chi_1, \chi_1; \chi_1) (R_1^d)^2 \chi_1^2 - 1/3 H_1^{F'}(1, \chi_m, \chi_{1s}, \chi_{1s}; \chi_1) R_{n1s}^m R_{1sm}^z \chi_1^2 \\
& - 2/3 H_1^{F'}(1, \chi_{1s}, \chi_{1s}, \chi_1; \chi_1) R_{n1s}^m R_{1sm}^z \chi_1^2 - 2/3 H_1^{F'}(3, \chi_{1s}, \chi_1, \chi_{1s}; \chi_1) R_1^d \chi_1^2 \\
& - 1/8 H_{21}^{F'}(1, \chi_m, \chi_{1s}, \chi_{1s}; \chi_1) R_{mn1s}^v \chi_1^2 + H_{21}^{F'}(1, \chi_1, \chi_{1s}, \chi_{1s}; \chi_1) [1/4 R_{s\pi\eta}^1 \chi_1^2 - 1/8 R_1^c \chi_1^2] \\
& + H_{21}^{F'}(1, \chi_s, \chi_{1s}, \chi_{1s}; \chi_1) [3/8 \chi_1^2 - 1/8 R_s^c \chi_1^2] + 3/4 H_{21}^{F'}(1, \chi_{st}, \chi_{1s}, \chi_{1t}; \chi_1) \chi_1^2 \\
& - 1/8 H_{21}^{F'}(2, \chi_1, \chi_{1s}, \chi_{1s}; \chi_1) R_1^d \chi_1^2. \tag{29}
\end{aligned}$$

B. Results for $d_{\text{val}} = 2$

In general, the expressions for $d_{\text{val}} = 2$ are longer than those for $d_{\text{val}} = 1$, as the number of independent integrals that can appear is significantly larger. The full result for $d_{\text{sea}} = 3$ at NNLO is very large, and has not yet been worked out. Consequently, the largest number of nondegenerate sea quarks considered for $d_{\text{val}} = 2$ is $d_{\text{sea}} = 2$. The NLO result for $d_{\text{val}} = 2$ and $d_{\text{sea}} = 2$, which agrees with the result of Ref. [5] is

$$\begin{aligned} \delta^{(4)22} = & 12 L_4^r \bar{\chi}_1 + 4 L_5^r \chi_{13} + \bar{A}(\chi_p) [1/6 R_{\eta\eta}^p \\ & - 1/12 R_p^c] + 1/4 \bar{A}(\chi_{ps}) - 1/12 \bar{A}(\chi_\eta) R_{\eta 13}^v \end{aligned}$$

$$- 1/12 \bar{B}(\chi_p, \chi_p, 0) R_p^d. \quad (30)$$

At NNLO, the contribution to the decay constant from the $\mathcal{O}(p^6)$ counterterms for $d_{\text{val}} = 2$ is similar to that for $d_{\text{val}} = 1$. For $d_{\text{sea}} = 2$, that contribution is

$$\begin{aligned} \delta_{\text{ct}}^{(6)22} = & 4 K_{19}^r \chi_p^2 + 24 K_{20}^r \bar{\chi}_1 \chi_{13} + 24 K_{21}^r \bar{\chi}_2 \\ & + 72 K_{22}^r \bar{\chi}_1^2 + 8 K_{23}^r \chi_1 \chi_3. \end{aligned} \quad (31)$$

The NNLO contributions to the decay constant from the chiral loops are, for $d_{\text{sea}} = 1$ and $d_{\text{sea}} = 2$, respectively,

$$\begin{aligned} \delta_{\text{loops}}^{(6)21} = & \pi_{16} L_0^r [-1/2 \chi_1 \chi_3 - 13/3 \chi_{13} \chi_4 + \chi_{13}^2 - 3/2 \chi_4^2] - 2 \pi_{16} L_1^r \chi_{13}^2 - \pi_{16} L_2^r [\chi_{13}^2 + 8 \chi_4^2] \\ & + \pi_{16} L_3^r [-7/12 \chi_1 \chi_3 - 17/6 \chi_{13} \chi_4 + 11/6 \chi_{13}^2 - 3/4 \chi_4^2] + \pi_{16}^2 [-59/384 \chi_1 \chi_3 - 1/2 \chi_{13} \chi_4 + 65/384 \chi_{13}^2 \\ & - 73/128 \chi_4^2] - 48 L_4^r L_5^r \chi_{13} \chi_4 - 72 L_4^r \chi_4^2 - 8 L_5^r \chi_{13}^2 + \bar{A}(\chi_p) \pi_{16} [5/96 \chi_p + 1/32 \chi_q + 1/48 \chi_4 \\ & - 1/12 R_q^p \chi_p - 1/8 R_q^p \chi_4] + \bar{A}(\chi_p) L_0^r [2/3 \chi_p + 8/3 R_q^p \chi_p + 2/3 R_p^d] + \bar{A}(\chi_p) L_3^r [5/3 \chi_p + 2/3 R_q^p \chi_p \\ & + 5/3 R_p^d] + \bar{A}(\chi_p) L_4^r [\chi_4 - 2 R_q^p \chi_4] + \bar{A}(\chi_p) L_5^r [1/6 \chi_p - 1/6 \chi_q - 1/3 \chi_4 - 2/3 R_q^p \chi_p] \\ & + \bar{A}(\chi_p)^2 [19/288 - 1/72 R_q^p R_p^d] - \bar{A}(\chi_p) \bar{A}(\chi_{p4}) [1/16 + 1/12 R_q^p] - \bar{A}(\chi_p) \bar{A}(\chi_{q4}) [3/16 + 1/12 R_q^p] \\ & + 1/8 \bar{A}(\chi_p) \bar{A}(\chi_{13}) + \bar{A}(\chi_p) \bar{B}(\chi_p, \chi_p; 0) [11/36 \chi_p - 1/18 R_q^p \chi_p - 1/72 R_q^p R_p^d + 1/144 R_p^d] \\ & + \bar{A}(\chi_p) \bar{B}(\chi_q, \chi_q; 0) [-1/72 R_q^p R_p^d + 1/144 R_q^d] - 1/4 \bar{A}(\chi_p) \bar{B}(\chi_{p4}, \chi_{p4}; 0) \chi_{p4} \\ & - 1/18 \bar{A}(\chi_p) \bar{B}(\chi_1, \chi_3; 0) R_p^q \chi_p + 1/18 \bar{A}(\chi_p) \bar{C}(\chi_p, \chi_p, \chi_p; 0) R_p^d \chi_p + 1/8 \bar{A}(\chi_p; \varepsilon) \pi_{16} [-\chi_p + R_q^p \chi_4] \\ & + \bar{A}(\chi_{p4}) \pi_{16} [3/16 \chi_{p4} - 9/16 \chi_{q4} - 9/16 \chi_4] - 6 \bar{A}(\chi_{p4}) L_0^r \chi_{p4} - 15 \bar{A}(\chi_{p4}) L_3^r \chi_{p4} - 9 \bar{A}(\chi_{p4}) L_4^r \chi_4 \\ & + 3 \bar{A}(\chi_{p4}) L_5^r \chi_{13} - 9/32 \bar{A}(\chi_{p4})^2 + \bar{A}(\chi_{p4}) \bar{B}(\chi_p, \chi_p; 0) [1/8 \chi_p - 5/8 \chi_{p4}] - 1/16 \bar{A}(\chi_{p4}) \bar{B}(\chi_q, \chi_q; 0) R_q^d \\ & + 1/6 \bar{A}(\chi_{p4}) \bar{B}(\chi_1, \chi_3; 0) \chi_4 + 1/3 \bar{A}(\chi_{p4}) \bar{B}(\chi_1, \chi_3; 0, k) + 9/8 \bar{A}(\chi_{p4}; \varepsilon) \pi_{16} \chi_4 + \bar{A}(\chi_1) \bar{A}(\chi_3) [-1/144 \\ & + 1/36 R_3^1 R_1^3] - 4 \bar{A}(\chi_{13}) L_1^r \chi_{13} - 10 \bar{A}(\chi_{13}) L_2^r \chi_{13} + 1/8 \bar{A}(\chi_{13})^2 - 1/2 \bar{A}(\chi_{13}) \bar{B}(\chi_1, \chi_3; 0, k) \\ & + 1/4 \bar{A}(\chi_{13}; \varepsilon) \pi_{16} \chi_{13} + 9/16 \bar{A}(\chi_{14}) \bar{A}(\chi_{34}) - 64 \bar{A}(\chi_4) L_1^r \chi_4 - 16 \bar{A}(\chi_4) L_2^r \chi_4 + 32 \bar{A}(\chi_4) L_4^r \chi_4 \\ & + 2/9 \bar{A}(\chi_4) \bar{B}(\chi_p, \chi_p; 0) \chi_4 - 4/9 \bar{A}(\chi_4) \bar{B}(\chi_1, \chi_3; 0) \chi_4 + \bar{A}(\chi_4; \varepsilon) \pi_{16} \chi_4 + \bar{B}(\chi_p, \chi_p; 0) \pi_{16} [1/96 R_p^d \chi_p \\ & + 1/32 R_p^d \chi_q + 1/16 R_p^d \chi_4] + 2/3 \bar{B}(\chi_p, \chi_p; 0) L_0^r R_p^d \chi_p + 5/3 \bar{B}(\chi_p, \chi_p; 0) L_3^r R_p^d \chi_p \\ & + \bar{B}(\chi_p, \chi_p; 0) L_4^r [2 \chi_p \chi_4 - 4 R_q^p \chi_p \chi_4 + 3 R_p^d \chi_4] + \bar{B}(\chi_p, \chi_p; 0) L_5^r [-2/3 \chi_p \chi_4 + 4/3 \chi_p^2 - 4/3 R_q^p \chi_p \chi_{13} \\ & - 1/3 R_p^d \chi_{13}] + \bar{B}(\chi_p, \chi_p; 0) L_6^r [4 \chi_4^2 - 8 R_q^p \chi_p \chi_4] + 4 \bar{B}(\chi_p, \chi_p; 0) L_7^r (R_p^d)^2 + \bar{B}(\chi_p, \chi_p; 0) L_8^r [4/3 \chi_4^2 \\ & - 8/3 R_p^d \chi_p^2] + \bar{B}(\chi_p, \chi_p; 0)^2 [1/18 R_p^q R_p^d \chi_p + 1/288 (R_p^d)^2] - 1/18 \bar{B}(\chi_p, \chi_p; 0) \bar{B}(\chi_1, \chi_3; 0) R_p^q R_p^d \chi_p \\ & + 1/18 \bar{B}(\chi_p, \chi_p; 0) \bar{C}(\chi_p, \chi_p, \chi_p; 0) (R_p^d)^2 \chi_p - 1/8 \bar{B}(\chi_p, \chi_p; 0, \varepsilon) \pi_{16} R_p^d \chi_{p4} \\ & - 18 \bar{B}(\chi_{p4}, \chi_{p4}; 0) L_4^r \chi_{p4} \chi_4 - 6 \bar{B}(\chi_{p4}, \chi_{p4}; 0) L_5^r \chi_{p4}^2 + 36 \bar{B}(\chi_{p4}, \chi_{p4}; 0) L_6^r \chi_{p4} \chi_4 \\ & + 12 \bar{B}(\chi_{p4}, \chi_{p4}; 0) L_8^r \chi_{p4}^2 + 1/144 \bar{B}(\chi_1, \chi_1; 0) \bar{B}(\chi_3, \chi_3; 0) R_1^d R_3^d - 8 \bar{B}(\chi_1, \chi_3; 0) L_7^r R_1^d R_3^d \\ & - 8/3 \bar{B}(\chi_1, \chi_3; 0) L_8^r R_1^d R_3^d + 4 \bar{C}(\chi_p, \chi_p, \chi_p; 0) L_4^r R_p^d \chi_p \chi_4 + 4/3 \bar{C}(\chi_p, \chi_p, \chi_p; 0) L_5^r R_p^d \chi_p^2 \\ & - 8 \bar{C}(\chi_p, \chi_p, \chi_p; 0) L_6^r R_p^d \chi_p \chi_4 - 8/3 \bar{C}(\chi_p, \chi_p, \chi_p; 0) L_8^r R_p^d \chi_p^2 + H^F(1, \chi_p, \chi_p, \chi_{13}; \chi_{13}) [1/8 \chi_p \\ & - 5/72 \chi_{13} + 1/36 R_q^p R_p^q \chi_{13}] + H^F(1, \chi_p, \chi_{14}, \chi_{34}; \chi_{13}) [-1/16 \chi_p + 1/16 \chi_q - 1/8 R_q^p \chi_4] \\ & + H^F(1, \chi_1, \chi_{13}, \chi_3; \chi_{13}) [1/72 \chi_{13} - 1/18 R_3^1 R_1^3 \chi_{13}] - 1/8 H^F(1, \chi_{13}, \chi_{13}, \chi_{13}; \chi_{13}) \chi_{13} \\ & - H^F(1, \chi_{14}, \chi_{34}, \chi_4; \chi_{13}) \chi_4 + H^F(2, \chi_p, \chi_p, \chi_{13}; \chi_{13}) [1/36 R_q^p R_p^d \chi_{13} - 1/72 R_p^d \chi_{13}] \\ & + H^F(2, \chi_p, \chi_{13}, \chi_q; \chi_{13}) [1/36 R_p^q R_p^d \chi_{13} - 1/72 R_p^d \chi_{13}] + 1/8 H^F(2, \chi_p, \chi_{14}, \chi_{34}; \chi_{13}) R_p^d \chi_{p4} \\ & - 1/144 H^F(5, \chi_p, \chi_p, \chi_{13}; \chi_{13}) (R_p^d)^2 \chi_{13} - 1/72 H^F(5, \chi_1, \chi_3, \chi_{13}; \chi_{13}) R_1^d R_3^d \chi_{13} \end{aligned}$$

$$\begin{aligned}
& + H^{F'}(1, \chi_p, \chi_p, \chi_{13}; \chi_{13}) [-1/8 \chi_p \chi_{13} - 5/24 \chi_{13}^2 + 1/12 R_q^p \chi_{13}^2 - 1/36 (R_q^p)^2 \chi_{13}^2] \\
& + H^{F'}(1, \chi_p, \chi_{14}, \chi_{34}; \chi_{13}) [3/16 \chi_p \chi_{13} - 1/16 \chi_{13} \chi_4 - 1/8 R_q^p R_p^d \chi_{13} + 3/8 R_p^q \chi_{13}^2] \\
& + H^{F'}(1, \chi_1, \chi_{13}, \chi_3; \chi_{13}) [-1/72 \chi_{13}^2 + 1/6 R_3^1 R_1^3 \chi_{13}^2] + 1/8 H^{F'}(1, \chi_{13}, \chi_{13}, \chi_{13}; \chi_{13}) \chi_{13}^2 \\
& + H^{F'}(1, \chi_{14}, \chi_{34}, \chi_4; \chi_{13}) \chi_{13} \chi_4 + H^{F'}(2, \chi_p, \chi_p, \chi_{13}; \chi_{13}) [-1/36 R_q^p R_p^d \chi_{13}^2 + 5/72 R_p^d \chi_{13}^2] \\
& + H^{F'}(2, \chi_p, \chi_{13}, \chi_q; \chi_{13}) [-1/36 R_p^q R_p^d \chi_{13}^2 + 1/72 R_p^d \chi_{13}^2] - 1/8 H^{F'}(2, \chi_p, \chi_{14}, \chi_{34}; \chi_{13}) R_p^d \chi_{p4} \chi_{13} \\
& + 5/144 H^{F'}(5, \chi_p, \chi_p, \chi_{13}; \chi_{13}) (R_p^d)^2 \chi_{13}^2 + 1/72 H^{F'}(5, \chi_1, \chi_3, \chi_{13}; \chi_{13}) R_1^d R_3^d \chi_{13}^2 \\
& + H_1^{F'}(1, \chi_p, \chi_p, \chi_{13}; \chi_{13}) [10/9 \chi_{13}^2 - 2/9 R_q^p R_p^q \chi_{13}^2] - H_1^{F'}(1, \chi_p, \chi_{14}, \chi_{34}; \chi_{13}) R_p^q \chi_{13}^2 \\
& - H_1^{F'}(1, \chi_{p4}, \chi_{q4}, \chi_p; \chi_{13}) \chi_{13}^2 + H_1^{F'}(1, \chi_{13}, \chi_1, \chi_3; \chi_{13}) [1/9 \chi_{13}^2 - 2/9 R_3^1 R_1^3 \chi_{13}^2] \\
& + H_1^{F'}(3, \chi_{13}, \chi_p, \chi_p; \chi_{13}) [1/9 R_q^p R_p^d \chi_{13}^2 - 1/9 R_p^d \chi_{13}^2] + 1/9 H_1^{F'}(3, \chi_{13}, \chi_p, \chi_q; \chi_{13}) R_p^q R_p^d \chi_{13}^2 \\
& - 1/18 H_1^{F'}(7, \chi_{13}, \chi_p, \chi_p; \chi_{13}) (R_p^d)^2 \chi_{13}^2 - 3/8 H_{21}^{F'}(1, \chi_p, \chi_p, \chi_{13}; \chi_{13}) \chi_{13}^2 \\
& + 3/8 H_{21}^{F'}(1, \chi_p, \chi_{14}, \chi_{34}; \chi_{13}) R_p^q \chi_{13}^2 + 3/8 H_{21}^{F'}(1, \chi_{p4}, \chi_p, \chi_{q4}; \chi_{13}) R_p^p \chi_{13}^2 \\
& - 3/8 H_{21}^{F'}(1, \chi_{p4}, \chi_q, \chi_{q4}; \chi_{13}) R_p^p \chi_{13}^2 + H_{21}^{F'}(1, \chi_{13}, \chi_p, \chi_p; \chi_{13}) [5/24 \chi_{13}^2 - 1/12 R_q^p R_p^q \chi_{13}^2] \\
& + H_{21}^{F'}(1, \chi_{13}, \chi_1, \chi_3; \chi_{13}) [-1/24 \chi_{13}^2 + 1/6 R_3^1 R_1^3 \chi_{13}^2] + 3/8 H_{21}^{F'}(1, \chi_{13}, \chi_{13}, \chi_{13}; \chi_{13}) \chi_{13}^2 \\
& + 3 H_{21}^{F'}(1, \chi_4, \chi_{14}, \chi_{34}; \chi_{13}) \chi_{13}^2 - 3/8 H_{21}^{F'}(3, \chi_{p4}, \chi_p, \chi_{q4}; \chi_{13}) R_p^d \chi_{13}^2 \\
& + H_{21}^{F'}(3, \chi_{13}, \chi_p, \chi_p; \chi_{13}) [-1/12 R_q^p R_p^d \chi_{13}^2 + 1/24 R_p^d \chi_{13}^2] + H_{21}^{F'}(3, \chi_{13}, \chi_p, \chi_q; \chi_{13}) [-1/12 R_p^q R_p^d \chi_{13}^2 \\
& + 1/24 R_p^d \chi_{13}^2] + 1/48 H_{21}^{F'}(7, \chi_{13}, \chi_p, \chi_p; \chi_{13}) (R_p^d)^2 \chi_{13}^2 + 1/24 H_{21}^{F'}(7, \chi_{13}, \chi_1, \chi_3; \chi_{13}) R_1^d R_3^d \chi_{13}^2, \quad (32)
\end{aligned}$$

$$\begin{aligned}
\delta_{\text{loops}}^{(6)22} & = \pi_{16} L_0^r [4/9 \chi_\eta \chi_4 - 1/2 \chi_1 \chi_3 + \chi_{13}^2 - 13/3 \bar{\chi}_1 \chi_{13} - 35/18 \bar{\chi}_2] - 2 \pi_{16} L_1^r \chi_{13}^2 \\
& - \pi_{16} L_2^r [11/3 \chi_\eta \chi_4 + \chi_{13}^2 + 13/3 \bar{\chi}_2] + \pi_{16} L_3^r [4/9 \chi_\eta \chi_4 - 7/12 \chi_1 \chi_3 + 11/6 \chi_{13}^2 - 17/6 \bar{\chi}_1 \chi_{13} - 43/36 \bar{\chi}_2] \\
& + \pi_{16}^2 [-15/64 \chi_\eta \chi_4 - 59/384 \chi_1 \chi_3 + 65/384 \chi_{13}^2 - 1/2 \bar{\chi}_1 \chi_{13} - 43/128 \bar{\chi}_2] - 48 L_4^r L_5^r \bar{\chi}_1 \chi_{13} - 72 L_4^r \bar{\chi}_1^2 \\
& - 8 L_5^r \chi_{13}^2 + \bar{A}(\chi_p) \pi_{16} [-1/24 \chi_p + 1/48 \bar{\chi}_1 - 1/8 \bar{\chi}_1 R_{q\eta}^p + 1/16 \bar{\chi}_1 R_p^c - 1/48 R_{q\eta}^p \chi_p - 1/16 R_{q\eta}^p \chi_q \\
& + 1/48 R_{pp}^p \chi_\eta + 1/16 R_p^c \chi_{13}] + \bar{A}(\chi_p) L_0^r [8/3 R_{q\eta}^p \chi_p + 2/3 R_p^c \chi_p + 2/3 R_p^d] + \bar{A}(\chi_p) L_3^r [2/3 R_{q\eta}^p \chi_p \\
& + 5/3 R_p^c \chi_p + 5/3 R_p^d] + \bar{A}(\chi_p) L_4^r [-2 \bar{\chi}_1 \bar{\chi}_{\eta\eta 0}^{pp} - 2 \bar{\chi}_1 R_{q\eta}^p + 3 \bar{\chi}_1 R_p^c] + \bar{A}(\chi_p) L_5^r [-2/3 \bar{\chi}_{\eta\eta 1}^{pp} - R_{q\eta}^p \chi_p \\
& + 1/3 R_{q\eta}^p \chi_q + 1/2 R_p^c \chi_p - 1/6 R_p^c \chi_q] + \bar{A}(\chi_p)^2 [1/16 + 1/72 (R_{q\eta}^p)^2 - 1/72 R_{q\eta}^p R_p^c + 1/288 (R_p^c)^2] \\
& + \bar{A}(\chi_p) \bar{A}(\chi_{ps}) [-1/36 R_{q\eta}^p - 5/72 R_{s\eta}^p + 7/144 R_p^c] - \bar{A}(\chi_p) \bar{A}(\chi_{qs}) [1/36 R_{q\eta}^p + 1/24 R_{s\eta}^p + 1/48 R_p^c] \\
& + \bar{A}(\chi_p) \bar{A}(\chi_\eta) [-1/72 R_{q\eta}^p R_{\eta 13}^p + 1/144 R_p^c R_{\eta 13}^p] + 1/8 \bar{A}(\chi_p) \bar{A}(\chi_{13}) + 1/12 \bar{A}(\chi_p) \bar{A}(\chi_{46}) R_{pp}^p \\
& + \bar{A}(\chi_p) \bar{B}(\chi_p, \chi_p; 0) [1/4 \chi_p - 1/18 R_{q\eta}^p R_p^c \chi_p - 1/72 R_{q\eta}^p R_p^d + 1/18 (R_p^c)^2 \chi_p + 1/144 R_p^c R_p^d] \\
& + \bar{A}(\chi_p) \bar{B}(\chi_p, \chi_\eta; 0) [1/18 R_{pp}^p R_p^c \chi_p - 1/18 R_{13}^p R_p^c \chi_p] + \bar{A}(\chi_p) \bar{B}(\chi_q, \chi_q; 0) [-1/72 R_{q\eta}^p R_q^d + 1/144 R_p^c R_q^d] \\
& - 1/12 \bar{A}(\chi_p) \bar{B}(\chi_{ps}, \chi_{ps}; 0) R_{s\eta}^p \chi_{ps} - 1/18 \bar{A}(\chi_p) \bar{B}(\chi_1, \chi_3; 0) R_{q\eta}^p R_p^c \chi_p \\
& + 1/18 \bar{A}(\chi_p) \bar{C}(\chi_p, \chi_p, \chi_p; 0) R_p^c R_p^d \chi_p + \bar{A}(\chi_p; \varepsilon) \pi_{16} [1/8 \bar{\chi}_1 R_{q\eta}^p - 1/16 \bar{\chi}_1 R_p^c - 1/16 R_p^c \chi_p - 1/16 R_p^d] \\
& + \bar{A}(\chi_{ps}) \pi_{16} [1/16 \chi_{ps} - 3/16 \chi_{qs} - 3/16 \bar{\chi}_1] - 2 \bar{A}(\chi_{ps}) L_0^r \chi_{ps} - 5 \bar{A}(\chi_{ps}) L_3^r \chi_{ps} - 3 \bar{A}(\chi_{ps}) L_4^r \bar{\chi}_1 \\
& + \bar{A}(\chi_{ps}) L_5^r \chi_{13} + \bar{A}(\chi_{ps}) \bar{A}(\chi_\eta) [7/144 R_{pp}^p - 5/72 R_{ps}^p - 1/48 R_{qq}^p + 5/72 R_{qs}^p - 1/36 R_{13}^p] \\
& + \bar{A}(\chi_{ps}) \bar{B}(\chi_p, \chi_p; 0) [1/24 R_{s\eta}^p \chi_p - 5/24 R_{s\eta}^p \chi_{ps}] + \bar{A}(\chi_{ps}) \bar{B}(\chi_p, \chi_\eta; 0) [-1/18 R_{ps}^p R_{qp\eta}^p \chi_p \\
& - 1/9 R_{ps}^p R_{qp\eta}^p \chi_{ps}] - 1/48 \bar{A}(\chi_{ps}) \bar{B}(\chi_q, \chi_q; 0) R_q^d + 1/18 \bar{A}(\chi_{ps}) \bar{B}(\chi_1, \chi_3; 0) R_{s\eta}^p \chi_s \\
& + 1/9 \bar{A}(\chi_{ps}) \bar{B}(\chi_1, \chi_3; 0, k) R_{s\eta}^p + 3/16 \bar{A}(\chi_{ps}; \varepsilon) \pi_{16} [\chi_s + \bar{\chi}_1] - 1/8 \bar{A}(\chi_{p4})^2 - 1/8 \bar{A}(\chi_{p4}) \bar{A}(\chi_{p6}) \\
& + 1/8 \bar{A}(\chi_{p4}) \bar{A}(\chi_{q6}) - 1/32 \bar{A}(\chi_{p6})^2 + \bar{A}(\chi_\eta) \pi_{16} [1/16 \bar{\chi}_1 R_{\eta 13}^p - 1/48 R_{\eta 13}^p \chi_\eta + 1/16 R_{\eta 13}^p \chi_{13}] \\
& + \bar{A}(\chi_\eta) L_0^r [4 R_{13}^p \chi_\eta + 2/3 R_{\eta 13}^p \chi_\eta] - 8 \bar{A}(\chi_\eta) L_1^r \chi_\eta - 2 \bar{A}(\chi_\eta) L_2^r \chi_\eta + \bar{A}(\chi_\eta) L_3^r [4 R_{13}^p \chi_\eta + 5/3 R_{\eta 13}^p \chi_\eta] \\
& + \bar{A}(\chi_\eta) L_4^r [4 \chi_\eta + \bar{\chi}_1 R_{\eta 13}^p] - \bar{A}(\chi_\eta) L_5^r [1/6 R_{pp}^p \chi_q + R_{13}^p \chi_{13} + 1/6 R_{\eta 13}^p \chi_\eta] + 1/288 \bar{A}(\chi_\eta)^2 (R_{\eta 13}^p)^2 \\
& + 1/12 \bar{A}(\chi_\eta) \bar{A}(\chi_{46}) R_{\eta 13}^p + \bar{A}(\chi_\eta) \bar{B}(\chi_p, \chi_p; 0) [-1/36 \bar{\chi}_{\eta\eta 1}^{pp} - 1/18 R_{q\eta}^p R_{pp}^p \chi_p + 1/18 R_{pp}^p R_p^c \chi_p \\
& + 1/144 R_p^d R_{\eta 13}^p] + \bar{A}(\chi_\eta) \bar{B}(\chi_p, \chi_\eta; 0) [-1/18 \bar{\chi}_{\eta\eta 1}^{pp} + 1/18 \bar{\chi}_{q\eta 1}^{pp} + 1/18 (R_{pp}^p)^2 R_{qp\eta}^p \chi_p]
\end{aligned}$$

$$\begin{aligned}
& - 1/12 \bar{A}(\chi_\eta) \bar{B}(\chi_{ps}, \chi_{ps}; 0) R_{ps}^\eta \chi_{ps} - \bar{A}(\chi_\eta) \bar{B}(\chi_\eta, \chi_\eta; 0) [1/216 R_{\eta 13}^v \chi_4 + 1/27 R_{\eta 13}^v \chi_6] \\
& - 1/18 \bar{A}(\chi_\eta) \bar{B}(\chi_1, \chi_3; 0) R_{\eta\eta}^1 R_{\eta\eta}^3 \chi_\eta + 1/18 \bar{A}(\chi_\eta) \bar{C}(\chi_p, \chi_p, \chi_p; 0) R_{pp}^\eta R_p^d \chi_p + \bar{A}(\chi_\eta; \varepsilon) \pi_{16} [1/8 \chi_\eta \\
& - 1/16 \bar{\chi}_1 R_{\eta 13}^v - 1/8 R_{13}^\eta \chi_\eta - 1/16 R_{\eta 13}^v \chi_\eta] + \bar{A}(\chi_1) \bar{A}(\chi_3) [-1/72 R_{q\eta}^p R_q^c + 1/36 R_{3\eta}^1 R_{1\eta}^3 + 1/144 R_1^c R_3^c] \\
& - 4 \bar{A}(\chi_{13}) L_1^r \chi_{13} - 10 \bar{A}(\chi_{13}) L_2^r \chi_{13} + 1/8 \bar{A}(\chi_{13})^2 - 1/2 \bar{A}(\chi_{13}) \bar{B}(\chi_1, \chi_3; 0, k) \\
& + 1/4 \bar{A}(\chi_{13}; \varepsilon) \pi_{16} \chi_{13} + 1/4 \bar{A}(\chi_{14}) \bar{A}(\chi_{34}) + 1/16 \bar{A}(\chi_{16}) \bar{A}(\chi_{36}) - 24 \bar{A}(\chi_4) L_1^r \chi_4 - 6 \bar{A}(\chi_4) L_2^r \chi_4 \\
& + 12 \bar{A}(\chi_4) L_4^r \chi_4 + 1/12 \bar{A}(\chi_4) \bar{B}(\chi_p, \chi_p; 0) (R_{4\eta}^p)^2 \chi_4 + 1/6 \bar{A}(\chi_4) \bar{B}(\chi_p, \chi_\eta; 0) [R_{4\eta}^p R_{p4}^\eta \chi_4 - R_{4\eta}^p R_{q4}^\eta \chi_4] \\
& - 1/24 \bar{A}(\chi_4) \bar{B}(\chi_\eta, \chi_\eta; 0) R_{\eta 13}^v \chi_4 - 1/6 \bar{A}(\chi_4) \bar{B}(\chi_1, \chi_3; 0) R_{4\eta}^1 R_{4\eta}^3 \chi_4 + 3/8 \bar{A}(\chi_4; \varepsilon) \pi_{16} \chi_4 \\
& - 32 \bar{A}(\chi_{46}) L_1^r \chi_{46} - 8 \bar{A}(\chi_{46}) L_2^r \chi_{46} + 16 \bar{A}(\chi_{46}) L_4^r \chi_{46} + \bar{A}(\chi_{46}) \bar{B}(\chi_p, \chi_p; 0) [1/9 \chi_{46} + 1/12 R_{pp}^\eta \chi_p \\
& + 1/36 R_{pp}^\eta \chi_4 + 1/9 R_{p4}^\eta \chi_6] + \bar{A}(\chi_{46}) \bar{B}(\chi_p, \chi_\eta; 0) [-1/18 R_{pp}^\eta \chi_4 - 1/9 R_{p4}^\eta \chi_6 + 1/9 R_{q4}^\eta \chi_6 + 1/18 R_{13}^\eta \chi_4] \\
& - 1/6 \bar{A}(\chi_{46}) \bar{B}(\chi_p, \chi_\eta; 0, k) [R_{pp}^\eta - R_{13}^\eta] + 1/9 \bar{A}(\chi_{46}) \bar{B}(\chi_\eta, \chi_\eta; 0) R_{\eta 13}^v \chi_{46} - \bar{A}(\chi_{46}) \bar{B}(\chi_1, \chi_3; 0) [2/9 \chi_{46} \\
& + 1/9 R_{p4}^\eta \chi_6 + 1/18 R_{13}^\eta \chi_4] - 1/6 \bar{A}(\chi_{46}) \bar{B}(\chi_1, \chi_3; 0, k) R_{13}^\eta + 1/2 \bar{A}(\chi_{46}; \varepsilon) \pi_{16} \chi_{46} \\
& + \bar{B}(\chi_p, \chi_p; 0) \pi_{16} [1/16 \bar{\chi}_1 R_p^d + 1/96 R_p^d \chi_p + 1/32 R_p^d \chi_q] + 2/3 \bar{B}(\chi_p, \chi_p; 0) L_0^r R_p^d \chi_p \\
& + 5/3 \bar{B}(\chi_p, \chi_p; 0) L_3^r R_p^d \chi_p + \bar{B}(\chi_p, \chi_p; 0) L_4^r [-2 \bar{\chi}_1 \bar{\chi}_{\eta\eta 0}^{pp} \chi_p - 4 \bar{\chi}_1 R_{q\eta}^p \chi_p + 4 \bar{\chi}_1 R_p^c \chi_p + 3 \bar{\chi}_1 R_p^d] \\
& + \bar{B}(\chi_p, \chi_p; 0) L_5^r [-2/3 \bar{\chi}_{\eta\eta 1}^{pp} \chi_p - 4/3 R_{q\eta}^p \chi_p^2 + 4/3 R_p^c \chi_p^2 + 1/2 R_p^d \chi_p - 1/6 R_p^d \chi_q] \\
& + \bar{B}(\chi_p, \chi_p; 0) L_6^r [4 \bar{\chi}_1 \bar{\chi}_{\eta\eta 1}^{pp} + 8 \bar{\chi}_1 R_{q\eta}^p \chi_p - 8 \bar{\chi}_1 R_p^c \chi_p] + 4 \bar{B}(\chi_p, \chi_p; 0) L_7^r (R_p^d)^2 \\
& + \bar{B}(\chi_p, \chi_p; 0) L_8^r [4/3 \bar{\chi}_{\eta\eta 2}^{pp} + 8/3 R_{q\eta}^p \chi_p^2 - 8/3 R_p^c \chi_p^2] + \bar{B}(\chi_p, \chi_p; 0)^2 [-1/18 R_{q\eta}^p R_p^d \chi_p + 1/18 R_p^c R_p^d \chi_p \\
& + 1/288 (R_p^d)^2] + 1/18 \bar{B}(\chi_p, \chi_p; 0) \bar{B}(\chi_p, \chi_\eta; 0) [R_{pp}^\eta R_p^d \chi_p - R_{13}^\eta R_p^d \chi_p] \\
& - 1/18 \bar{B}(\chi_p, \chi_p; 0) \bar{B}(\chi_1, \chi_3; 0) R_{p\eta}^q R_p^d \chi_p + 1/18 \bar{B}(\chi_p, \chi_p; 0) \bar{C}(\chi_p, \chi_p, \chi_p; 0) (R_p^d)^2 \chi_p \\
& - 1/16 \bar{B}(\chi_p, \chi_p; 0, \varepsilon) \pi_{16} [\bar{\chi}_1 R_p^d + R_p^d \chi_p] + 8 \bar{B}(\chi_p, \chi_\eta; 0) L_7^r R_{qp\eta}^z R_p^d R_{\eta 46p}^z \\
& + 8/3 \bar{B}(\chi_p, \chi_\eta; 0) L_8^r R_{qp\eta}^z R_p^d R_{\eta 46p}^z - 6 \bar{B}(\chi_{ps}, \chi_{ps}; 0) L_4^r \bar{\chi}_1 \chi_{ps} - 2 \bar{B}(\chi_{ps}, \chi_{ps}; 0) L_5^r \chi_{ps}^2 \\
& + 12 \bar{B}(\chi_{ps}, \chi_{ps}; 0) L_6^r \bar{\chi}_1 \chi_{ps} + 4 \bar{B}(\chi_{ps}, \chi_{ps}; 0) L_8^r \chi_{ps}^2 + 2 \bar{B}(\chi_\eta, \chi_\eta; 0) L_4^r \bar{\chi}_1 R_{\eta 13}^v \chi_\eta \\
& + 2/3 \bar{B}(\chi_\eta, \chi_\eta; 0) L_5^r R_{\eta 13}^v \chi_\eta^2 - 4 \bar{B}(\chi_\eta, \chi_\eta; 0) L_6^r \bar{\chi}_1 R_{\eta 13}^v \chi_\eta + 4 \bar{B}(\chi_\eta, \chi_\eta; 0) L_7^r R_{311\eta\eta}^z (R_{\eta 461}^z)^2 \\
& - \bar{B}(\chi_\eta, \chi_\eta; 0) L_8^r [4/9 R_{\eta 13}^v \chi_4^2 + 8/9 R_{\eta 13}^v \chi_6^2] + 1/144 \bar{B}(\chi_1, \chi_1; 0) \bar{B}(\chi_3, \chi_3; 0) R_1^d R_3^d \\
& - 8 \bar{B}(\chi_1, \chi_3; 0) L_7^r R_1^d R_3^d - 8/3 \bar{B}(\chi_1, \chi_3; 0) L_8^r R_1^d R_3^d + 4 \bar{C}(\chi_p, \chi_p, \chi_p; 0) L_4^r \bar{\chi}_1 R_p^d \chi_p \\
& + 4/3 \bar{C}(\chi_p, \chi_p, \chi_p; 0) L_5^r R_p^d \chi_p^2 - 8 \bar{C}(\chi_p, \chi_p, \chi_p; 0) L_6^r \bar{\chi}_1 R_p^d \chi_p - 8/3 \bar{C}(\chi_p, \chi_p, \chi_p; 0) L_8^r R_p^d \chi_p^2 \\
& + H^F(1, \chi_p, \chi_p, \chi_{13}; \chi_{13}) [1/8 \chi_p - 1/16 \chi_{13} - 1/36 (R_{q\eta}^p)^2 \chi_{13} + 1/36 R_{q\eta}^p R_p^c \chi_{13} - 1/144 (R_p^c)^2 \chi_{13}] \\
& + H^F(1, \chi_p, \chi_{1s}, \chi_{3s}; \chi_{13}) [-1/12 R_{q\eta}^p \chi_{qs} + 1/24 R_{q\eta}^p \chi_{13} - 1/16 R_{s\eta}^p \chi_p + 1/48 R_{s\eta}^p \chi_q + 1/24 R_p^c \chi_{ps}] \\
& - 1/4 H^F(1, \chi_{p4}, \chi_{q6}, \chi_{46}; \chi_{13}) \chi_{46} + H^F(1, \chi_\eta, \chi_p, \chi_{13}; \chi_{13}) [1/36 R_{q\eta}^p R_{\eta 13}^v \chi_{13} - 1/72 R_p^c R_{\eta 13}^v \chi_{13}] \\
& - 1/144 H^F(1, \chi_\eta, \chi_\eta, \chi_{13}; \chi_{13}) (R_{\eta 13}^v)^2 \chi_{13} - 1/48 H^F(1, \chi_\eta, \chi_{1s}, \chi_{3s}; \chi_{13}) [R_{ps}^\eta \chi_p - R_{ps}^\eta \chi_q \\
& - R_{\eta ps}^v \chi_\eta - R_{\eta 13}^v \chi_s] + H^F(1, \chi_1, \chi_{13}, \chi_3; \chi_{13}) [1/36 R_{q\eta}^p R_q^c \chi_{13} - 1/18 R_{3\eta}^1 R_{1\eta}^3 \chi_{13} \\
& - 1/72 R_1^c R_3^c \chi_{13}] - 1/8 H^F(1, \chi_{13}, \chi_{13}, \chi_{13}; \chi_{13}) \chi_{13} - 3/8 H^F(1, \chi_{14}, \chi_{34}, \chi_4; \chi_{13}) \chi_4 \\
& + H^F(2, \chi_p, \chi_p, \chi_{13}; \chi_{13}) [1/36 R_{q\eta}^p R_p^d \chi_{13} - 1/72 R_p^c R_p^d \chi_{13}] - 1/72 H^F(2, \chi_p, \chi_\eta, \chi_{13}; \chi_{13}) R_p^d R_{\eta 13}^v \chi_{13} \\
& + H^F(2, \chi_p, \chi_{13}, \chi_q; \chi_{13}) [1/36 R_{p\eta}^q R_p^d \chi_{13} - 1/72 R_q^c R_p^d \chi_{13}] + 1/24 H^F(2, \chi_p, \chi_{1s}, \chi_{3s}; \chi_{13}) R_p^d \chi_{ps} \\
& - 1/144 H^F(5, \chi_p, \chi_p, \chi_{13}; \chi_{13}) (R_p^d)^2 \chi_{13} - 1/72 H^F(5, \chi_1, \chi_3, \chi_{13}; \chi_{13}) R_1^d R_3^d \chi_{13} \\
& + H^{F'}(1, \chi_p, \chi_{1s}, \chi_{3s}; \chi_{13}) [-5/24 R_{q\eta}^p \chi_q \chi_{13} + 1/12 R_{q\eta}^p \chi_{ps} \chi_{13} + 1/16 R_{s\eta}^p \chi_q \chi_{13} + 7/48 R_{s\eta}^p \chi_{13} \chi_s \\
& - 1/24 R_p^c \chi_{ps} \chi_{13}] + 1/4 H^{F'}(1, \chi_{p4}, \chi_{q6}, \chi_{46}; \chi_{13}) \chi_{13} \chi_{46} + H^{F'}(1, \chi_\eta, \chi_p, \chi_{13}; \chi_{13}) [1/9 R_{q\eta}^p R_{\eta 13}^v \chi_{13}^2 \\
& - 1/36 R_{q\eta}^p R_{\eta 13}^v \chi_{13}^2 + 1/18 R_{pp}^\eta R_p^c \chi_{13}^2 + 1/72 R_p^c R_{\eta 13}^v \chi_{13}^2] + H^{F'}(1, \chi_\eta, \chi_\eta, \chi_{13}; \chi_{13}) [1/9 (R_{13}^\eta)^2 \chi_{13}^2 \\
& + 1/9 R_{13}^\eta R_{\eta 13}^v \chi_{13}^2 + 5/144 (R_{\eta 13}^v)^2 \chi_{13}^2] + H^{F'}(1, \chi_\eta, \chi_{1s}, \chi_{3s}; \chi_{13}) [-1/16 R_{ps}^\eta \chi_p \chi_{13} - 5/48 R_{ps}^\eta \chi_q \chi_{13} \\
& + 1/6 R_{1s}^\eta R_{\eta s}^z \chi_{13}^2 - 1/48 R_{\eta ps}^v \chi_\eta \chi_{13} - 1/48 R_{\eta 13}^v \chi_{13} \chi_s] + H^{F'}(1, \chi_1, \chi_{13}, \chi_3; \chi_{13}) [-1/36 R_{q\eta}^p R_q^c \chi_{13}^2 \\
& + 1/6 R_{3\eta}^1 R_{1\eta}^3 \chi_{13}^2 + 1/72 R_1^c R_3^c \chi_{13}^2] + H^{F'}(1, \chi_{13}, \chi_p, \chi_p; \chi_{13}) [-1/8 \chi_p \chi_{13} - 3/16 \chi_{13}^2 - 1/36 (R_{q\eta}^p)^2 \chi_{13}^2]
\end{aligned}$$

$$\begin{aligned}
& + 1/12 R_{q\eta}^p R_p^c \chi_{13}^2 - 1/48 (R_p^c)^2 \chi_{13}^2] + 1/8 H^{F'}(1, \chi_{13}, \chi_{13}, \chi_{13}; \chi_{13}) \chi_{13}^2 \\
& + 3/8 H^{F'}(1, \chi_{14}, \chi_{34}, \chi_4; \chi_{13}) \chi_{13} \chi_4 + H^{F'}(2, \chi_p, \chi_p, \chi_{13}; \chi_{13}) [-1/36 R_{q\eta}^p R_p^d \chi_{13}^2 + 5/72 R_p^c R_p^d \chi_{13}^2] \\
& + H^{F'}(2, \chi_p, \chi_\eta, \chi_{13}; \chi_{13}) [1/18 R_{pp}^\eta R_p^d \chi_{13}^2 + 1/72 R_p^d R_{\eta 13}^v \chi_{13}^2] \\
& + H^{F'}(2, \chi_p, \chi_{13}, \chi_q; \chi_{13}) [-1/36 R_{p\eta}^q R_p^d \chi_{13}^2 + 1/72 R_q^c R_p^d \chi_{13}^2] - 1/24 H^{F'}(2, \chi_p, \chi_{1s}, \chi_{3s}; \chi_{13}) R_p^d \chi_{ps} \chi_{13} \\
& + 5/144 H^{F'}(5, \chi_p, \chi_p, \chi_{13}; \chi_{13}) (R_p^d)^2 \chi_{13}^2 + 1/72 H^{F'}(5, \chi_1, \chi_3, \chi_{13}; \chi_{13}) R_1^d R_3^d \chi_{13}^2 \\
& + H_1^{F'}(1, \chi_p, \chi_p, \chi_{13}; \chi_{13}) [\chi_{13}^2 + 2/9 (R_{q\eta}^p)^2 \chi_{13}^2 - 2/9 R_{q\eta}^p R_p^c \chi_{13}^2 + 1/9 (R_p^c)^2 \chi_{13}^2] \\
& + 1/3 H_1^{F'}(1, \chi_p, \chi_{1s}, \chi_{3s}; \chi_{13}) R_{q\eta}^p R_{sqp}^z \chi_{13}^2 - 1/3 H_1^{F'}(1, \chi_{ps}, \chi_{qs}, \chi_p; \chi_{13}) R_{s\eta}^p \chi_{13}^2 \\
& - 1/3 H_1^{F'}(1, \chi_{ps}, \chi_{qs}, \chi_\eta; \chi_{13}) R_{13}^\eta R_{spq}^z \chi_{13}^2 + 1/9 H_1^{F'}(1, \chi_{13}, \chi_p, \chi_\eta; \chi_{13}) [R_{p\eta}^p R_{\eta 13}^v \chi_{13}^2 \\
& - R_{pp}^\eta R_{qp\eta}^z R_p^c \chi_{13}^2] - H_1^{F'}(1, \chi_{13}, \chi_\eta, \chi_\eta; \chi_{13}) [1/9 R_{13}^\eta R_{\eta 13}^v \chi_{13}^2 + 1/18 (R_{\eta 13}^v)^2 \chi_{13}^2] \\
& + H_1^{F'}(1, \chi_{13}, \chi_1, \chi_3; \chi_{13}) [1/9 R_{q\eta}^p R_q^c \chi_{13}^2 - 2/9 R_{3\eta}^1 R_{1\eta}^3 \chi_{13}^2] + 1/9 H_1^{F'}(3, \chi_{13}, \chi_p, \chi_p; \chi_{13}) [R_{q\eta}^p R_p^d \chi_{13}^2 \\
& - R_p^c R_p^d \chi_{13}^2] + 1/9 H_1^{F'}(3, \chi_{13}, \chi_p, \chi_q; \chi_{13}) R_{p\eta}^q R_p^d \chi_{13}^2 + 1/9 H_1^{F'}(3, \chi_{13}, \chi_p, \chi_\eta; \chi_{13}) R_{13}^p R_{pq\eta}^z R_p^d \chi_{13}^2 \\
& - 1/18 H_1^{F'}(7, \chi_{13}, \chi_p, \chi_p; \chi_{13}) (R_p^d)^2 \chi_{13}^2 - 3/8 H_{21}^{F'}(1, \chi_p, \chi_p, \chi_{13}; \chi_{13}) \chi_{13}^2 \\
& - 1/8 H_{21}^{F'}(1, \chi_p, \chi_{1s}, \chi_{3s}; \chi_{13}) R_{q\eta}^p R_{sqp}^z \chi_{13}^2 + 1/8 H_{21}^{F'}(1, \chi_{ps}, \chi_{qs}, \chi_p; \chi_{13}) [R_{q\eta}^p \chi_{13}^2 + R_{s\eta}^p \chi_{13}^2 - R_p^c \chi_{13}^2] \\
& + 1/8 H_{21}^{F'}(1, \chi_{ps}, \chi_{qs}, \chi_q; \chi_{13}) R_{p\eta}^q R_{spq}^z \chi_{13}^2 + 1/8 H_{21}^{F'}(1, \chi_{ps}, \chi_{qs}, \chi_\eta; \chi_{13}) R_{13}^\eta R_{pq\eta}^z R_{sp\eta}^z \chi_{13}^2 \\
& - 1/8 H_{21}^{F'}(1, \chi_\eta, \chi_{1s}, \chi_{3s}; \chi_{13}) R_{13}^\eta R_{s1\eta}^z R_{s3\eta}^z \chi_{13}^2 + H_{21}^{F'}(1, \chi_{13}, \chi_p, \chi_p; \chi_{13}) [3/16 \chi_{13}^2 + 1/12 (R_{q\eta}^p)^2 \chi_{13}^2 \\
& - 1/12 R_{q\eta}^p R_p^c \chi_{13}^2 + 1/48 (R_p^c)^2 \chi_{13}^2] + H_{21}^{F'}(1, \chi_{13}, \chi_p, \chi_\eta; \chi_{13}) [-1/12 R_{q\eta}^p R_{\eta 13}^v \chi_{13}^2 + 1/24 R_p^c R_{\eta 13}^v \chi_{13}^2] \\
& + 1/48 H_{21}^{F'}(1, \chi_{13}, \chi_\eta, \chi_\eta; \chi_{13}) (R_{\eta 13}^v)^2 \chi_{13}^2 + H_{21}^{F'}(1, \chi_{13}, \chi_1, \chi_3; \chi_{13}) [-1/12 R_{q\eta}^p R_q^c \chi_{13}^2 + 1/6 R_{3\eta}^1 R_{1\eta}^3 \chi_{13}^2 \\
& + 1/24 R_1^c R_3^c \chi_{13}^2] + 3/8 H_{21}^{F'}(1, \chi_{13}, \chi_{13}, \chi_{13}; \chi_{13}) \chi_{13}^2 + 9/8 H_{21}^{F'}(1, \chi_4, \chi_{14}, \chi_{34}; \chi_{13}) \chi_{13}^2 \\
& + 3/4 H_{21}^{F'}(1, \chi_{46}, \chi_{p4}, \chi_{q6}; \chi_{13}) \chi_{13}^2 - 1/8 H_{21}^{F'}(3, \chi_{ps}, \chi_p, \chi_{qs}; \chi_{13}) R_p^d \chi_{13}^2 \\
& + H_{21}^{F'}(3, \chi_{13}, \chi_p, \chi_p; \chi_{13}) [-1/12 R_{q\eta}^p R_p^d \chi_{13}^2 + 1/24 R_p^c R_p^d \chi_{13}^2] + H_{21}^{F'}(3, \chi_{13}, \chi_p, \chi_q; \chi_{13}) [-1/12 R_{p\eta}^q R_p^d \chi_{13}^2 \\
& + 1/24 R_q^c R_p^d \chi_{13}^2] + 1/24 H_{21}^{F'}(3, \chi_{13}, \chi_p, \chi_\eta; \chi_{13}) R_p^d R_{\eta 13}^v \chi_{13}^2 + 1/48 H_{21}^{F'}(7, \chi_{13}, \chi_p, \chi_p; \chi_{13}) (R_p^d)^2 \chi_{13}^2 \\
& + 1/24 H_{21}^{F'}(7, \chi_{13}, \chi_1, \chi_3; \chi_{13}) R_1^d R_3^d \chi_{13}^2.
\end{aligned} \tag{33}$$

IV. DISCUSSION AND CONCLUSIONS

The analytical formulas for the NNLO contributions to the decay constants are very complicated, especially when the valence quark masses are nondegenerate. The practical usefulness of those results thus depends completely on the availability of convenient numerical implementations. In the near future, we will make such an implementation available [14].

The results, as given in the preceding section, depend on nine $\mathcal{O}(p^4)$ and five $\mathcal{O}(p^6)$ LEC:s. Eventually, their proper values should be determined from lattice QCD by a fit of the NLO + NNLO formulas to the simulation data. At the present time, these input parameters have been taken from the continuum work in NNLO χ PT of Ref. [15]. The combination of LEC:s used in this paper corresponds to 'fit 10' of Ref. [15], which had $F_0 = 87.7$ MeV and $\mu = 770$ MeV. The parameters which were not determined by that fit, namely L_0^r, L_4^r and L_6^r , have been set to zero, as have all the K_i^r . This has been done at a scale of $\mu = 770$ MeV. It should be noted that L_0^r cannot be determined from experiment, but is obtainable from

partially quenched simulations. Some recent results on L_4^r and L_6^r may be found in Ref. [16].

The graphical presentation of multidimensional functions is a more immediate problem. In the most general case, the decay constant of a charged pseudoscalar meson in PQ χ PT is a function of two valence and three sea quark masses. The plots presented in this section should therefore be viewed as an attempt to cover the potentially most interesting parts of the parameter space. They also serve as a consistency check for the formulas in the preceding section, as the cases with fewer different masses should be obtained numerically as limits of the more general cases.

A. Numerical Results for $d_{\text{sea}} = 1$

To this end, the result for $d_{\text{val}} = 1$ and $d_{\text{sea}} = 1$ has been plotted along different rays in the χ_1 - χ_4 plane, characterized by an angle θ which is defined according to

$$\tan(\theta) = \chi_4/\chi_1, \tag{34}$$

such that $\theta = 45^\circ$ corresponds to the unquenched case of equal sea and valence quark masses. The typical situation in lattice QCD simulations, where sea quarks are heavier than their valence counterparts, is attained for $\theta > 45^\circ$. The shift in the decay constant due to the combined NLO + NNLO contributions has been plotted in Fig. 2 for different values of θ . The quantity plotted, Δ , denotes the relative change in the decay constant and is defined in accordance with Eq. (24), as

$$F_{\text{phys}} = F_0 (1 + \Delta). \quad (35)$$

Thus, when plotted along a curve of the form (34), Δ is expected to vanish in the chiral limit, which is indeed borne out in all plots presented in this section. This shows that the NNLO shifts to the decay constant in PQ χ PT do not produce unphysical logarithms. On the other hand, the curve labeled 'A' in Fig. 2 is plotted for a constant sea mass with $\chi_4 = 0.125 \text{ GeV}^2$, and in that case Δ approaches a constant as $\chi_1 \rightarrow 0$.




FIG. 2: The combined NLO and NNLO shifts of the decay constant, plotted for $d_{\text{val}} = 1$ and $d_{\text{sea}} = 1$, for values of θ between 15° and 75° . The curve 'A' corresponds to $\chi_4 = 0.125 \text{ GeV}^2$. In order to reduce clutter, most of the lines in this plot, with the exception of the 15° line, have not been drawn all the way to the origin.

The results along the lines (34) at NLO is shown in Fig. 3. The chiral expansion for the decay constants converges better than for the masses as can be seen by comparison of Figs. 2 and 3 with the corresponding ones in Ref. [6]. Note, however, that the dependence of Δ on the numerical values of the LEC:s is strong and can change the behavior of the NNLO results quite considerably. It should also be kept in mind that many of the curves plotted in Fig. 2 reach far beyond the expected radius of convergence of PQ χ PT.




FIG. 3: The NLO shift of the decay constant, plotted for $d_{\text{val}} = 1$ and $d_{\text{sea}} = 1$, for values of θ between 15° and 75° . The curve 'A' corresponds to $\chi_4 = 0.125 \text{ GeV}^2$.




FIG. 4: The combined NLO and NNLO shifts of the decay constant, plotted for $d_{\text{val}} = 1$, $d_{\text{sea}} = 2$, $\theta = 60^\circ$, and a strange sea quark mass parameter z between $1/2$ and 10 .

B. Numerical Results for $d_{\text{sea}} = 2$

The physically more interesting case of $d_{\text{sea}} = 2$, where the strange sea quark mass may deviate from that of the u, d quarks, can then be accounted for by the introduction of an additional parameter z , which is defined as

$$z = \chi_6 / \chi_4. \quad (36)$$

The change in Δ when the strange sea quark mass is allowed to vary is shown in Fig. 4. In general, the NNLO effects appear to become larger as the strange sea quark mass is increased. Fig. 4 also shows that Eq. (28) reduces numerically to the $d_{\text{val}} = 1$, $d_{\text{sea}} = 1$ result of Eq. (27) when all sea quark masses become equal, i.e. $z \rightarrow 1$.




FIG. 5: The combined NLO and NNLO shifts of the decay constant, plotted for $d_{\text{val}} = 2$, $d_{\text{sea}} = 2$, $z = 5$ and $\theta = 60^\circ$. The strange valence quark mass parameter x varies between 1 and 2.

It is also instructive to investigate the NNLO correction to the decay constant for $d_{\text{val}} = 2$ and $d_{\text{sea}} = 2$. The behavior of Δ when the strange valence quark mass deviates from that of the u, d quarks is shown in Fig. 5, where the results have been parameterized in terms of

$$x = \chi_3 / \chi_1. \quad (37)$$

It is evident that Δ changes rapidly with increasing x for values of χ_1 above about 0.15 GeV^2 . Also, Eq. (33) is seen to reduce correctly to the $d_{\text{val}} = 1$, $d_{\text{sea}} = 2$ result when $x \rightarrow 1$.

C. Numerical Results for $d_{\text{sea}} = 3$

In the case of $d_{\text{sea}} = 3$, χ_π and χ_η are non-trivially related to the sea quark masses, and consequently the sea quark sector is more complicated. The variation in the mass of the d quark with respect to the u quark, in the sea quark sector, has been parameterized by

$$y = \chi_5 / \chi_4. \quad (38)$$

Sample plots of Δ for $d_{\text{val}} = 1$, with all three sea quark masses different, are shown in Fig. 6. Those plots also demonstrate numerically the consistency of Eq. (29) with the $d_{\text{sea}} = 1, 2$ results.




FIG. 6: The combined NLO and NNLO shifts of the decay constant, plotted for $d_{\text{val}} = 1$, $d_{\text{sea}} = 3$, $\theta = 60^\circ$ and $z = 5$. The mass parameter y of the down quark in the sea sector ranges between 1 and 2.

D. Conclusions

A generic feature of the plots presented in this section is that the curves for Δ show a pronounced dip around $\chi_1 = 0.1 \text{ GeV}^2$. This indicates a strong cancellation between the NLO and NNLO contributions to Δ . It should be noted that such a feature is apparently not exhibited by the analogous expressions for the pseudoscalar meson mass [6]. This cancellation also depends strongly on the choice of the LEC:s. Many more plots can of course be produced but those presented give a first indication of the size of the corrections and how they vary with the different quark masses used as input. We have not attempted any fit of our results to the available lattice data; Many simulations are performed with only two flavors of sea quarks, and in addition we need extrapolations to zero lattice spacing and infinite volume at each quark mass to apply the present formulas. Work for the case with two sea quark flavors is in progress.

In conclusion, we have calculated the decay constants of the charged, or off-diagonal, pseudoscalar mesons to NNLO in PQ χ PT and presented analytical as well as numerical results for a variety of different combinations of quark masses. The NNLO contributions were found, as expected from previous work in NNLO χ PT [13, 15, 17], to be sizable even though there is a tendency toward cancellation with the NLO result. As the results depend on a number of largely unknown LEC:s, statements about the convergence of the chiral expansion have to be postponed at this time. The NNLO effects are definitely non-negligible at presently used quark masses in Lattice QCD simulations.

Acknowledgments

The program **FORM 3.0** has been used extensively in these calculations [18]. This work is supported by the European Union TMR network, Contract No. HPRN-CT-2002-00311 (EURIDICE). TL wishes to thank the

Thomas Jefferson National Accelerator Facility (TJNAF) and the Helsinki Institute of Physics (HIP), where part of this paper was completed, for their hospitality. TL also thanks the Mikael Björnberg memorial foundation for a travel grant.

-
- [1] S. Weinberg, *Physica A* **96** (1979) 327; J. Gasser and H. Leutwyler, *Ann. Phys.* **158** (1984) 142; J. Gasser and H. Leutwyler, *Nucl. Phys. B* **250**, 465 (1985).
 - [2] C. W. Bernard and M. F. L. Golterman, *Phys. Rev. D* **46**, 853 (1992) [arXiv:hep-lat/9204007].
 - [3] C. W. Bernard and M. F. L. Golterman, *Phys. Rev. D* **49**, 486 (1994) [arXiv:hep-lat/9306005].
 - [4] S. R. Sharpe and N. Shoresh, *Phys. Rev. D* **62**, 094503 (2000) [arXiv:hep-lat/0006017].
 - [5] S. R. Sharpe and N. Shoresh, *Phys. Rev. D* **64**, 114510 (2001) [arXiv:hep-lat/0108003].
 - [6] J. Bijnens, N. Danielsson and T. A. Lähde, *Phys. Rev. D* **70**, 111503 (2004) [arXiv:hep-lat/0406017].
 - [7] F. Farchioni, I. Montvay and E. Scholz [qq+q Collaboration], *Eur. Phys. J. C* **37**, 197 (2004) [arXiv:hep-lat/0403014].
 - [8] C. Aubin *et al.* [MILC Collaboration], *Phys. Rev. D* **70**, 114501 (2004) [arXiv:hep-lat/0407028].
 - [9] J. Bijnens, N. Danielsson and T. A. Lähde, to be published.
 - [10] P. H. Damgaard and K. Splittorff, *Phys. Rev. D* **62**, 054509 (2000) [arXiv:hep-lat/0003017].
 - [11] J. Bijnens, G. Colangelo and G. Ecker, *JHEP* **9902**, 020 (1999) [arXiv:hep-ph/9902437].
 - [12] J. Bijnens, G. Colangelo and G. Ecker, *Ann. Phys.* **280**, 100 (2000) [arXiv:hep-ph/9907333].
 - [13] G. Amorós, J. Bijnens and P. Talavera, *Nucl. Phys. B* **568**, 319 (2000) [arXiv:hep-ph/9907264].
 - [14] The analytical formulas are available via <http://www.thep.lu.se/~bijnens/chpt.html>. The programs are available from the authors.
 - [15] G. Amorós, J. Bijnens and P. Talavera, *Nucl. Phys. B* **602**, 87 (2001) [arXiv:hep-ph/0101127].
 - [16] J. Bijnens, P. Dhonte and P. Talavera, *JHEP* **0401**, 050 (2004) [arXiv:hep-ph/0401039], *JHEP* **0405**, 036 (2004) [arXiv:hep-ph/0404150].
 - [17] E. Golowich and J. Kambor, *Phys. Rev. D* **58**, 036004 (1998) [arXiv:hep-ph/9710214].
 - [18] J. A. Vermaseren, arXiv:math-ph/0010025.